



# 1. Introduction

---

The following notes aim to provide a very informal introduction to Stochastic Calculus, and especially to the Itô integral. They owe a great deal to Dan Crisan's *Stochastic Calculus and Applications* lectures of 1998; and also much to various books especially those of L. C. G. Rogers and D. Williams, and Dellacherie and Meyer. They have also benefitted from lecture notes of T. Kurtz. They aren't supposed to be a set of lecture notes; and it's important to remember this if you are reading this as a part III student with a view to being examined in this course. Many important parts of the course have been omitted completely.

The present notes grew out of a set of typed notes which I produced when revising for the Part III course; combining the printed notes and my own handwritten notes into a consistent text. I've subsequently expanded them inserting some extra proofs. The notes principally concentrate on the parts of the course which I found hard; thus there is often little or no comment on more standard matters; as a secondary goal they aim to present the results in a form which can be readily extended :x their evolution they have taken a very informal style; in some ways I hope this may make them easier to read.

The goal of the notes in their current form is to present a fairly clear approach to the Itô integral with respect to *continuous* semimartingales. The various alternative approaches to this subject which can be found in books tend to divide into those presenting the integral directed entirely at Brownian Motion, and those who wish to prove results in complete generality for a semimartingale. Here at all points clarity has hopefully been the main goal here, rather than completeness; although secretly the approach aims to be readily extended to the discontinuous theory. I'd especially like to convince the reader that the Itô integral isn't that much harder in concept than the Lebesgue Integral with which we are all familiar. The motivating principle is to try and explain every detail, no matter how trivial it may seem once the subject has been understood!

Passages enclosed in boxes are intended to be viewed as digressions from the main text; usually describing an alternative approach, or giving an informal description of what is going on – feel free to skip these sections if you find them unhelpful. Some of the proof also contain solutions to important questions on the example sheets. I've tried to indicate this by the legend QS.

These notes contain errors with probability one. I always welcome people telling me about the errors because then I can fix them! I can be readily contacted by email as [afrb2@statslab.cam.ac.uk](mailto:afrb2@statslab.cam.ac.uk). Also suggestions for improvements or other additions are welcome.

Alan Bain

## 2. Contents

---

<b>1. Introduction</b>	<i>i</i>
<b>2. Contents</b>	<i>ii</i>
<b>3. Stochastic Processes</b>	1
3.1. Probability Space	1
3.2. Stochastic Process	1
<b>4. Martingales</b>	3
4.1. Stopping Times	3
<b>5. Basics</b>	6
5.1. Local Martingales	6
<b>6. Total Variation and the Stieltjes Integral</b>	9
6.1. Why we need a Stochastic Integral	9
6.2. Previsibility	10
6.3. Lebesgue-Stieltjes Integral	10
<b>7. The Integral</b>	13
7.1. Elementary Processes	13
7.2. Strictly Simple and Simple Processes	13
<b>8. The Stochastic Integral</b>	15
8.1. Integral for $H \in \mathcal{L}$ and $M \in \mathcal{M}_2$	15
8.2. Quadratic Variation	17
8.3. Covariation	20
8.4. Extension of the Integral to $L^2(M)$	21
8.5. Localisation	24
8.6. Some Important Results	25
<b>9. Semimartingales</b>	27
<b>10. Relations to Sums</b>	29
10.1. The UCP topology	29
10.2. Approximation via Riemann Sums	30
<b>11. Itô's Formula</b>	33
11.1. Applications of Itô's Formula	38
11.2. Exponential Martingales	39
<b>12. Lévy Characterisation of Brownian Motion</b>	41
<b>13. Time Change of Brownian Motion</b>	43
13.1. Gaussian Martingales	44
<b>14. Girsanov's Theorem</b>	46
14.1. Change of measure	46
<b>15. Relations to Second Order PDEs</b>	48
15.1. Infinitesimal Generator	48
15.2. The Dirichlet Problem	49
15.3. The Cauchy Problem	51
15.4. Feynman-Kač Representation	53
<b>16. Stochastic Filtering</b>	55

16.1. Signal Process . . . . .	55
16.2. Observation Process . . . . .	55
16.3. The Filtering Problem . . . . .	55
16.4. Change of Measure . . . . .	56
16.5. The Unnormalised Conditional Distribution . . . . .	61
16.6. The Zakai Equation . . . . .	62
16.7. Kushner-Stratonowich Equation . . . . .	67

### 3. Stochastic Processes

---

The following notes are a summary of results from the theory of stochastic processes, proofs may be found in the usual books, and in some cases in the notes from Dr. Norris' advanced probability course.

#### 3.1. Probability Space

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space. The set of  $\mathbb{P}$ -null subsets of  $\Omega$  is defined by

$$\mathcal{N} := \{N \subset \Omega : N \subset A \text{ for } A \in \mathcal{F}, \text{ with } \mathbb{P}(A) = 0\}.$$

The space is *complete* if for  $A \subset B \subset \Omega$  with  $B \in \mathcal{F}$  and  $\mathbb{P}(B) = 0$  then this implies that  $A \in \mathcal{F}$ .

In addition to the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , let  $(E, \mathcal{E})$  be a measurable space, the state space, which is typically  $(\mathbb{R}, \mathcal{B})$ , or  $(\mathbb{R}^n, \mathcal{B})$ . A random variable is a  $\mathcal{F}/\mathcal{E}$  measurable function  $X : \Omega \rightarrow E$ .

#### 3.2. Stochastic Process

Given a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and a measurable state space  $(E, \mathcal{E})$ , a stochastic process is a family  $(X_t)_{t \geq 0}$  such that  $X_t$  is an  $E$  valued random variable for each time  $t \geq 0$ .

##### Definition 1. Measurable Process

The process  $(X_t)_{t \geq 0}$  is said to be measurable if the mapping  $(t, \omega) \mapsto X_t(\omega)$  is measurable on  $\mathbb{R} \times \Omega$  with respect to the product  $\sigma$ -field  $\mathcal{B}(\mathbb{R}) \times \mathcal{F}$ .

Associated with a process is a *filtration*, an increasing chain of  $\sigma$ -algebras i.e.

$$\mathcal{F}_s \subset \mathcal{F}_t \text{ if } 0 \leq s \leq t < \infty.$$

Define  $\mathcal{F}_\infty$  by

$$\mathcal{F}_\infty = \sigma \left( \bigcup_{t \geq 0} \mathcal{F}_t \right).$$

If  $(X_t)_{t \geq 0}$  is a stochastic process, then the *natural filtration* of  $(X_t)_{t \geq 0}$  is given by

$$\mathcal{F}_t^X := \sigma(X_s : s \leq t).$$

The process  $(X_t)_{t \geq 0}$  is said to be  $(\mathcal{F}_t)_{t \geq 0}$  *adapted*, if  $X_t$  is  $\mathcal{F}_t$  measurable for each  $t \geq 0$ . The process  $(X_t)_{t \geq 0}$  is obviously adapted with respect to the natural filtration.

##### Definition 3.2.

A process  $(X_t)_{t \geq 0}$  is said to be *bounded* if there exists a universal constant  $K$  such that for all  $\omega$  and  $t \geq 0$ , then  $|X_t(\omega)| < K$ .

##### Definition 3.3.

Let  $X = (X_t)_{t \geq 0}$  be a stochastic process defined on  $(\Omega, \mathcal{F}, \mathbb{P})$ , and let  $X' = (X'_t)_{t \geq 0}$  be a stochastic process defined on  $(\Omega, \mathcal{F}, \mathbb{P})$ . Then  $X$  and  $X'$  have the same finite dimensional distributions if for all  $n$ ,  $0 \leq t_1 < t_2 < \dots < t_n < \infty$ , and  $A_1, A_2, \dots, A_n \in \mathcal{E}$ ,

$$\mathbb{P}(X_{t_1} \in A_1, X_{t_2} \in A_2, \dots, X_{t_n} \in A_n) = \mathbb{P}'(X'_{t_1} \in A_1, X'_{t_2} \in A_2, \dots, X'_{t_n} \in A_n).$$

**Definition 3.4.**

Let  $X$  and  $X'$  be defined on  $(\Omega, \mathcal{F}, \mathbb{P})$ . Then  $X$  and  $X'$  are modifications of each other if and only if

$$\mathbb{P}(\{\omega \in \Omega : X_t(\omega) = X'_t(\omega)\}) = 1 \quad \forall t \geq 0.$$

**Definition 3.5.**

Let  $X$  and  $X'$  be defined on  $(\Omega, \mathcal{F}, \mathbb{P})$ . Then  $X$  and  $X'$  are indistinguishable if and only if

$$\mathbb{P}(\{\omega \in \Omega : X_t(\omega) = X'_t(\omega) \forall t \geq 0\}) = 1.$$

There is a chain of implications

$$\text{indistinguishable} \Rightarrow \text{modifications} \Rightarrow \text{same f.d.d.}$$

## 4. Martingales

---

### Definition 4.1.

Let  $X = \{X_t, \mathcal{F}_t, t \geq 0\}$  be an integrable process then  $X$  is a

- (i) **Martingale** if and only if  $\mathbb{E}(X_t | \mathcal{F}_s) = X_s$  a.s. for  $0 \leq s \leq t < \infty$
- (ii) **Supermartingale** if and only if  $\mathbb{E}(X_t | \mathcal{F}_s) \leq X_s$  a.s. for  $0 \leq s \leq t < \infty$
- (iii) **Submartingale** if and only if  $\mathbb{E}(X_t | \mathcal{F}_s) \geq X_s$  a.s. for  $0 \leq s \leq t < \infty$

### Theorem (Kolmogorov) 4.2.

Let  $X = \{X_t, \mathcal{F}_t, t \geq 0\}$  be an integrable process. Then define  $\mathcal{F}_{t+} := \bigcap_{\epsilon > 0} \mathcal{F}_{t+\epsilon}$  and also the partial augmentation of  $\mathcal{F}$  by  $\tilde{\mathcal{F}}_t = \sigma(\mathcal{F}_{t+}, \mathcal{N})$ . Then there exists a martingale  $\tilde{X} = \{\tilde{X}_t, \tilde{\mathcal{F}}_t, t \geq 0\}$  right continuous, with left limits (CADLAG) such that  $X$  and  $\tilde{X}$  are modifications of each other.

### Definition 4.3.

A martingale  $X = \{X_t, \mathcal{F}_t, t \geq 0\}$  is said to be an  $L^2$ -martingale or a square integrable martingale if  $\mathbb{E}(X_t^2) < \infty$  for every  $t \geq 0$ .

### Definition 4.4.

A process  $X = \{X_t, \mathcal{F}_t, t \geq 0\}$  is said to be  $L^p$  bounded if and only if  $\sup_{t \geq 0} \mathbb{E}(|X_t|^p) < \infty$ . The space of  $L^2$  bounded martingales is denoted by  $\mathcal{M}_2$ , and the subspace of continuous  $L^2$  bounded martingales is denoted  $\mathcal{M}_2^c$ .

### Definition 4.5.

A process  $X = \{X_t, \mathcal{F}_t, t \geq 0\}$  is said to be uniformly integrable if and only if

$$\sup_{t \geq 0} \mathbb{E}(|X_t| 1_{|X_t| \geq N}) \rightarrow 0 \text{ as } N \rightarrow \infty.$$

### Orthogonality of Martingale Increments

A frequently used property of a martingale  $M$  is the *orthogonality of increments property* which states that for a square integrable martingale  $M$ , and  $Y \in \mathcal{F}_s$  with  $\mathbb{E}(Y^2) < \infty$  then

$$\mathbb{E}[Y(M_t - M_s)] = 0 \quad \text{for } t \geq s.$$

*Proof*

Via Cauchy Schwartz inequality  $\mathbb{E}|Y(M_t - M_s)| < \infty$ , and so

$$\mathbb{E}(Y(M_t - M_s)) = \mathbb{E}(\mathbb{E}(Y(M_t - M_s) | \mathcal{F}_s)) = \mathbb{E}(Y \mathbb{E}(M_t - M_s | \mathcal{F}_s)) = 0.$$

□

A typical example is  $Y = M_s$ , whence  $\mathbb{E}(M_s(M_t - M_s)) = 0$  is obtained. A common application is to the difference of two squares, let  $t \geq s$  then

$$\begin{aligned} \mathbb{E}((M_t - M_s)^2 | \mathcal{F}_s) &= \mathbb{E}(M_t^2 | \mathcal{F}_s) - 2M_s \mathbb{E}(M_t | \mathcal{F}_s) + M_s^2 \\ &= \mathbb{E}(M_t^2 - M_s^2 | \mathcal{F}_s) = \mathbb{E}(M_t^2 | \mathcal{F}_s) - M_s^2. \end{aligned}$$

### 4.1. Stopping Times

A random variable  $T : \Omega \rightarrow [0, \infty)$  is a stopping time if and only if  $\{\omega : T(\omega) \leq t\} \in \mathcal{F}_t$ . Given a stochastic process  $X = (X_t)_{t \geq 0}$ , a stopped process  $X^T$  may be defined by

$$X^T(\omega) := X_{T(\omega) \wedge t}(\omega),$$

$$\mathcal{F}_t := \{A \in \mathcal{F} : A \cap \{T \leq t\} \in \mathcal{F}_t\}.$$

#### Theorem (Optional Stopping).

Let  $X$  be a right continuous integrable,  $\mathcal{F}_t$  adapted process. Then the following are equivalent:

- (i)  $X$  is a martingale.
- (ii)  $X^T$  is a martingale for all stopping times  $T$ .
- (iii)  $\mathbb{E}(X_T) = \mathbb{E}(X_0)$  for all bounded stopping times  $T$ .
- (iv)  $\mathbb{E}(X_T | \mathcal{F}_S) = X_S$  for all bounded stopping times  $S$  and  $T$  such that  $S \leq T$ . If in addition,  $X$  is uniformly integrable then (iv) holds for all stopping times (not necessarily bounded).

#### Theorem (Doob's Martingale Inequalities).

Let  $M = \{M_t, \mathcal{F}_t, t \geq 0\}$  be a uniformly integrable martingale, and let  $M^* := \sup_{t \geq 0} |M_t|$ . Then

- (i) Maximal Inequality. For  $\lambda > 0$ ,

$$\lambda \mathbb{P}(M^* \geq \lambda) \leq \mathbb{E}[|M_\infty| 1_{M^* < \infty}].$$

- (ii)  $L^p$  maximal inequality. For  $1 < p < \infty$ ,

$$\|M^*\|_p \leq \frac{p}{p-1} \|M_\infty\|_p.$$

Note that the norm used in stating the Doob  $L^p$  inequality is defined by

$$\|M\|_p = [\mathbb{E}(|M|^p)]^{1/p}.$$

#### Theorem (Martingale Convergence).

Let  $M = \{M_t, \mathcal{F}_t, t \geq 0\}$  be a martingale.

- (i) If  $M$  is  $L^p$  bounded then  $M_\infty(\omega) := \lim_{t \rightarrow \infty} M_t(\omega)$   $P$ -a.s.
- (ii) If moreover  $M$  is uniformly integrable then  $\lim_{t \rightarrow \infty} M_t(\omega) = M_\infty(\omega)$  in  $L^1$ . Then for all  $A \in L^1(\mathcal{F}_\infty)$ , there exists a martingale  $A_t$  such that  $\lim_{t \rightarrow \infty} A_t = A$ , and  $A_t = \mathbb{E}(A | \mathcal{F}_t)$ . Here  $\mathcal{F}_\infty := \lim_{t \rightarrow \infty} \mathcal{F}_t$ .
- (iii) If moreover  $M$  is  $L^p$  bounded then  $\lim_{t \rightarrow \infty} M_t = M_\infty$  in  $L^p$ , and for all  $A \in L^p(\mathcal{F}_\infty)$ , there exists a martingale  $A_t$  such that  $\lim_{t \rightarrow \infty} A_t = A$ , and  $A_t = \mathbb{E}(A | \mathcal{F}_t)$ .

#### Definition 4.6.

Let  $\mathcal{M}_2$  denote the set of  $L^2$ -bounded CADLAG martingales i.e. martingales  $M$  such that

$$\sup_{t \geq 0} M_t^2 < \infty.$$

Let  $\mathcal{M}_2^c$  denote the set of  $L^2$ -bounded CADLAG martingales which are continuous. A norm may be defined on the space  $\mathcal{M}_2$  by  $\|M\|^2 = \|M_\infty\|_2^2 = \mathbb{E}(M_\infty^2)$ .

From the conditional Jensen's inequality, since  $f(x) = x^2$  is convex,

$$\begin{aligned}\mathbb{E}(M_\infty^2 | \mathcal{F}_t) &\geq (\mathbb{E}(M_\infty | \mathcal{F}_t))^2 \\ \mathbb{E}(M_\infty^2 | \mathcal{F}_t) &\geq (\mathbb{E}M_t)^2.\end{aligned}$$

Hence taking expectations

$$\mathbb{E}M_t^2 \leq \mathbb{E}M_\infty^2,$$

and since by martingale convergence in  $L^2$ , we get  $\mathbb{E}(M_t^2) \rightarrow \mathbb{E}(M_\infty^2)$ , it is clear that

$$\mathbb{E}(M_\infty^2) = \sup_{t \geq 0} \mathbb{E}(M_t^2).$$

**Theorem 4.7.**

The space  $(\mathcal{M}_2, \|\cdot\|)$  (up to equivalence classes defined by modifications) is a Hilbert space, with  $\mathcal{M}_2^c$  a closed subspace.

*Proof*

We prove this by showing a one to one correspondence between  $\mathcal{M}_2$  (the space of square integrable martingales) and  $L^2(\mathcal{F}_\infty)$ . The bijection is obtained via

$$\begin{aligned}f : \mathcal{M}_2 &\rightarrow L^2(\mathcal{F}_\infty) \\ f : (M_t)_{t \geq 0} &\mapsto M_\infty \equiv \lim_{t \rightarrow \infty} M_t \\ g : L^2(\mathcal{F}_\infty) &\rightarrow \mathcal{M}_2 \\ g : M_\infty &\mapsto M_t \equiv \mathbb{E}(M_\infty | \mathcal{F}_t)\end{aligned}$$

Notice that

$$\sup_t \mathbb{E}M_t^2 = \|M_\infty\|_2^2 = \mathbb{E}(M_\infty^2) < \infty,$$

as  $M_t$  is a square integrable martingale. As  $L^2(\mathcal{F}_\infty)$  is a Hilbert space,  $\mathcal{M}_2$  inherits this structure.

To see that  $\mathcal{M}_2^c$  is a closed subspace of  $\mathcal{M}_2$ , consider a Cauchy sequence  $\{M^{(n)}\}$  in  $\mathcal{M}_2$ , equivalently  $\{M_\infty^{(n)}\}$  is Cauchy in  $L^2(\mathcal{F}_\infty)$ . Hence  $M_\infty^{(n)}$  converges to a limit,  $M_\infty$  say, in  $L^2(\mathcal{F}_\infty)$ . Let  $M_t := \mathbb{E}(M_\infty | \mathcal{F}_t)$ , then

$$\sup_{t \geq 0} |M_t^{(n)} - M_t| \rightarrow 0, \text{ in } L^2,$$

that is  $M^{(n)} \rightarrow M$  uniformly in  $L^2$ . Hence there exists a subsequence  $n(k)$  such that  $M^{n(k)} \rightarrow M$  uniformly; as a uniform limit of continuous functions is continuous,  $M \in \mathcal{M}_2^c$ . Thus  $\mathcal{M}_2^c$  is a closed subspace of  $\mathcal{M}$ .

## 5. Basics

---

### 5.1. Local Martingales

A martingale has already been defined, but a weaker definition will prove useful for stochastic calculus. Note that I'll often drop references to the filtration  $\mathcal{F}_t$ , but this nevertheless forms an essential part of the (local) martingale.

Just before we dive in and define a Local Martingale, maybe we should pause and consider the reason for considering them. The important property of local martingales will only be seen later in the notes; and as we frequently see in this subject it is one of *stability* that is, they are a class of objects which are closed under an operation, in this case under the stochastic integral – an integral of a previsible process with a local martingale integrator is a local martingale.

**Definition 5.1.**

$M = \{M_t, \mathcal{F}_t, 0 \leq t \leq \infty\}$  is a local martingale if and only if there exists a sequence of stopping times  $T_n$  tending to infinity such that  $M^{T_n}$  are martingales for all  $n$ . The space of local martingales is denoted  $\mathcal{M}_{loc}$ , and the subspace of continuous local martingales is denoted  $\mathcal{M}_{loc}^c$ .

Recall that a martingale  $(X_t)_{t \geq 0}$  is said to be bounded if there exists a universal constant  $K$  such that for all  $\omega$  and  $t \geq 0$ , then  $|X_t(\omega)| < K$ .

**Theorem (QS1 Q1) 5.2.**

Every bounded local martingale is a martingale.

*Proof*

Let  $T_n$  be a sequence of stopping times as in the definition of a local martingale. This sequence tends to infinity, so pointwise  $X_t^{T_n}(\omega) \rightarrow X_t(\omega)$ . Using the conditional form of the dominated convergence theorem (using the constant bound as the dominating function), for  $t \geq s \geq 0$

$$\lim_{n \rightarrow \infty} \mathbb{E}(X_t^{T_n} | \mathcal{F}_s) = \mathbb{E}(X_t | \mathcal{F}_s).$$

But as  $X^{T_n}$  is a (genuine) martingale,  $\mathbb{E}(X_t^{T_n} | \mathcal{F}_s) = X_s^{T_n} = X_{T_n \wedge s}$ ; so

$$\mathbb{E}(X_t | \mathcal{F}_s) = \lim_{n \rightarrow \infty} \mathbb{E}(X_t^{T_n} | \mathcal{F}_s) = \lim_{n \rightarrow \infty} X_s^{T_n} = X_s.$$

Hence  $X_t$  is a genuine martingale. □

**Proposition 5.3.**

The following are equivalent

- (i)  $M = \{M_t, \mathcal{F}_t, 0 \leq t \leq \infty\}$  is a continuous martingale.
- (ii)  $M = \{M_t, \mathcal{F}_t, 0 \leq t \leq \infty\}$  is a continuous local martingale and for all  $t \geq 0$ , the set  $\{M_T : T \text{ a stopping time, } T \leq t\}$  is uniformly integrable.

*Proof*

(i)  $\Rightarrow$  (ii) By optional stopping theorem, if  $T \leq t$  then  $M_T = \mathbb{E}(M_t | \mathcal{F}_T)$  hence the set is uniformly integrable.

### Local Martingales which are not Martingales

There do exist local martingales which are not themselves martingales. The following is an example. Let  $B_t$  be a  $d$  dimensional Brownian Motion starting from  $x$ . It can be shown using Itô's formula that a harmonic function of a Brownian motion is a local martingale (this is on the example sheet). From standard PDE theory it is known that for  $d \geq 3$ , the function

$$f(x) = \frac{1}{|x|^{d-2}}$$

is a harmonic function, hence  $X_t = 1/|B_t|^{d-2}$  is a local martingale. Now consider the  $L^p$  norm of this local martingale

$$\mathbb{E}_x |X_t|^p = \int \frac{1}{(2\pi t)^{d/2}} \exp\left(-\frac{|y-x|^2}{2t}\right) |y|^{-(d-2)p} dy.$$

Consider when this integral converges. There are no divergence problems for  $|y|$  large, the potential problem lies in the vicinity of the origin. Here the term

$$\frac{1}{(2\pi t)^{d/2}} \exp\left(-\frac{|y-x|^2}{2t}\right)$$

is bounded, so we only need to consider the remainder of the integrand integrated over a ball of unit radius about the origin which is bounded by

$$C \int_{B(0,1)} |y|^{-(d-2)p} dy,$$

for some constant  $C$ , which on transformation into polar co-ordinates yields a bound of the form

$$C' \int_0^1 r^{-(d-2)p} r^{d-1} dr,$$

with  $C'$  another constant. This is finite if and only if  $-(d-2)p + (d-1) > -1$  (standard integrals of the form  $1/r^k$ ). This in turn requires that  $p < d/(d-2)$ . So clearly  $\mathbb{E}_x |X_t|$  will be finite for all  $d \geq 3$ .

Now although  $\mathbb{E}_x |X_t| < \infty$  and  $X_t$  is a local martingale, we shall show that it is not a martingale. Note that  $(B_t - x)$  has the same distribution as  $\sqrt{t}(B_1 - x)$  under  $\mathbb{P}_x$  (the probability measure induced by the BM starting from  $x$ ). So as  $t \rightarrow \infty$ ,  $|B_t| \rightarrow \infty$  in probability and  $X_t \rightarrow 0$  in probability. As  $X_t \geq 0$ , we see that  $\mathbb{E}_x(X_t) = \mathbb{E}_x |X_t| < \infty$ . Now note that for any  $R < \infty$ , we can construct a bound

$$\mathbb{E}_x X_t \leq \frac{1}{(2\pi t)^{d/2}} \int_{|y| \leq R} |y|^{-(d-2)} dy + R^{-(d-2)},$$

which converges, and hence

$$\limsup_{t \rightarrow \infty} \mathbb{E}_x X_t \leq R^{-(d-2)}.$$

As  $R$  was chosen arbitrarily we see that  $\mathbb{E}_x X_t \rightarrow 0$ . But  $\mathbb{E}_x X_0 = |x|^{-(d-2)} > 0$ , which implies that  $\mathbb{E}_x X_t$  is not constant, and hence  $X_t$  is not a martingale.

(ii)  $\Rightarrow$  (i) It is required to prove that  $\mathbb{E}(M_0) = \mathbb{E}(M_T)$  for any bounded stopping time  $T$ . Then by local martingale property for any  $n$ ,

$$\mathbb{E}(M_0) = \mathbb{E}(M_{T \wedge T_n}),$$

uniform integrability then implies that

$$\lim_{n \rightarrow \infty} \mathbb{E}(M_{T \wedge T_n}) = \mathbb{E}(M_T).$$

## 6. Total Variation and the Stieltjes Integral

---

Let  $A : [0, \infty) \rightarrow \mathbb{R}$  be a CADLAG (continuous to right, with left limits) process. Let a *partition*  $\Pi = \{t_0, t_1, \dots, t_m\}$  have  $0 = t_0 \leq t_1 \leq \dots \leq t_m = t$ ; the mesh of the partition is defined by

$$\delta(\Pi) = \max_{1 \leq k \leq m} |t_k - t_{k-1}|.$$

The variation of  $A$  is then defined as the increasing process  $V$  given by,

$$V_t := \sup_{\Pi} \left\{ \sum_{k=1}^{n(\Pi)} |A_{t_k \wedge t} - A_{t_{k-1} \wedge t}| : 0 = t_0 \leq t_1 \leq \dots \leq t_n = t \right\}.$$

An alternative definition is given by

$$V_t^0 := \lim_{n \rightarrow \infty} \sum_1^{\infty} |A_{k2^{-n} \wedge t} - A_{(k-1)2^{-n} \wedge t}|.$$

These can be shown to be equivalent (for CADLAG processes), since trivially (use the dyadic partition),  $V_t^0 \leq V_t$ . It is also possible to show that  $V_t^0 \geq V_t$  for the total variation of a CADLAG process.

### Definition 6.1.

A process  $A$  is said to have *finite variation* if the associated variation process  $V$  is finite (i.e. if for every  $t$  and every  $\omega$ ,  $|V_t(\omega)| < \infty$ ).

### 6.1. Why we need a Stochastic Integral

Before delving into the depths of the integral it's worth stepping back for a moment to see why the 'ordinary' integral cannot be used on a path at a time basis (i.e. separately for each  $\omega \in \Omega$ ). Suppose we were to do this i.e. set

$$I_t(X) = \int_0^t X_s(\omega) dM_s(\omega),$$

for  $M \in \mathcal{M}_2^c$ ; but for an interesting martingale (i.e. one which isn't zero a.s.), the total variation is not finite, even on a bounded interval like  $[0, T]$ . Thus the Lebesgue-Stieltjes integral definition isn't valid in this case. To generalise we shall see that the quadratic variation is actually the 'right' variation to use (higher variations turn out to be zero and lower ones infinite – see QS2 Q2). But to start, we shall consider integrating a previsible process  $H_t$  with an integrator which is an increasing finite variation process. First we shall prove that a continuous local martingale of finite variation is zero.

### Proposition 6.2.

If  $M$  is a continuous local martingale of finite variation, starting from zero then  $M$  is identically zero.

*Proof*

Let  $V$  be the variation process of  $M$ . This  $V$  is a continuous, adapted process. Now define a sequence of stopping times  $S_n$  as the first time  $V$  exceeds  $n$ , i.e.  $S_n := \inf_t \{t \geq 0 : V_t \geq n\}$ . Then the martingale  $M^{S_n}$  is of bounded variation. It therefore suffices to prove the result for a bounded, continuous martingale  $M$  of bounded variation.

Fix  $t \geq 0$  and let  $\{0 = t_0, t_1, \dots, t_N = t\}$  be a partition of  $[0, t]$ . Then since  $M_0 = 0$  it is clear that,  $M_t^2 = \sum_{k=1}^N (M_{t_k}^2 - M_{t_{k-1}}^2)$ . Then via orthogonality of martingale increments

$$\begin{aligned} \mathbb{E}(M_t^2) &= \mathbb{E} \left( \sum_{k=1}^N (M_{t_k} - M_{t_{k-1}})^2 \right) \\ &\leq \mathbb{E} \left( V_t \sup_k |M_{t_k} - M_{t_{k-1}}| \right) \end{aligned}$$

The integrand is bounded by  $n^2$  (from definition of the stopping time  $S_n$ ), hence the expectation converges to zero as the modulus of the partition tends to zero by the bounded convergence theorem. Hence  $M \equiv 0$ .  $\square$

## 6.2. Previsibility

The term previsible has crept into the discussion earlier. Now is the time for a proper definition.

### Definition 6.3.

The previsible (or predictable)  $\sigma$ -field  $\mathcal{P}$  is the  $\sigma$ -field on  $\mathbb{R}^+ \times \Omega$  generated by the processes  $(X_t)_{t \geq 0}$ , adapted to  $\mathcal{F}_t$ , with left continuous paths on  $(0, \infty)$ .

### Remark

The same  $\sigma$ -field is generated by left continuous, right limits processes (i.e. càglàd processes) which are adapted to  $\mathcal{F}_{t-}$ , or indeed continuous processes  $(X_t)_{t \geq 0}$ , which are adapted to  $\mathcal{F}_{t-}$ .

### Definition 6.4.

A process  $(X_t)_{t \geq 0}$  is said to be previsible, if the mapping  $(t, \omega) \mapsto X_t(\omega)$  is measurable with respect to the previsible  $\sigma$ -field  $\mathcal{P}$ .

## 6.3. Lebesgue-Stieltjes Integral

[In the lecture notes for this course, the Lebesgue-Stieltjes integral is considered first for functions  $A$  and  $H$ ; here I consider processes on a pathwise basis.]

Let  $A$  be an increasing cadlag process. This induces a Borel measure  $dA$  on  $(0, \infty)$  such that

$$dA((s, t])(\omega) = A_t(\omega) - A_s(\omega).$$

Let  $H$  be a previsible process (as defined above). The Lebesgue-Stieltjes integral of  $H$  is defined with respect to an increasing process  $A$  by

$$(H \cdot A)_t(\omega) = \int_0^t H_s(\omega) dA_s(\omega),$$

whenever  $H \geq 0$  or  $(|H| \cdot A)_t < \infty$ .

As a notational aside, we shall write

$$(H \cdot A)_t \equiv \int_0^t H dX,$$

and later on we shall use

$$d(H \cdot X) \equiv H dX.$$

This definition may be extended to integrator of finite variation which are not increasing, by decomposing the process  $A$  of finite variation into a difference of two increasing processes, so  $A = A^+ - A^-$ , where  $A^\pm = (V \pm A)/2$  (here  $V$  is the total variation process for  $A$ ). The integral of  $H$  with respect to the finite variation process  $A$  is then defined by

$$(H \cdot A)_t(\omega) := (H \cdot A^+)_t(\omega) - (H \cdot A^-)_t(\omega),$$

whenever  $(|H| \cdot V)_t < \infty$ .

There are no really new concepts of the integral in the foregoing; it is basically the Lebesgue-Stieltjes integral extended from functions  $H(t)$  to processes in a pathwise fashion (that's why  $\omega$  has been included in those definitions as a reminder).

**Theorem (QS1 Q3) 6.5.**

If  $X$  is a non-negative continuous local martingale and  $\mathbb{E}(X_0) < \infty$  then  $X_t$  is a supermartingale. If additionally  $X$  has constant mean, i.e.  $\mathbb{E}(X_t) = \mathbb{E}(X_0)$  for all  $t$  then  $X_t$  is a martingale.

*Proof*

As  $X_t$  is a continuous local martingale there is a sequence of stopping times  $T_n \uparrow \infty$  such that  $X^{T_n}$  is a genuine martingale. From this martingale property

$$\mathbb{E}(X_t^{T_n} | \mathcal{F}_s) = X_s^{T_n}.$$

As  $X_t \geq 0$  we can apply the conditional form of Fatou's lemma, so

$$\mathbb{E}(X_t | \mathcal{F}_s) = \mathbb{E}(\liminf_{n \rightarrow \infty} X_t^{T_n} | \mathcal{F}_s) \leq \liminf_{n \rightarrow \infty} \mathbb{E}(X_t^{T_n} | \mathcal{F}_s) = \liminf_{n \rightarrow \infty} X_s^{T_n} = X_s.$$

Hence  $\mathbb{E}(X_t | \mathcal{F}_s) \leq X_s$ , so  $X_t$  is a supermartingale.

Given the constant mean property  $\mathbb{E}(X_t) = \mathbb{E}(X_s)$ . Let

$$A_n := \{\omega : X_s - \mathbb{E}(X_t | \mathcal{F}_s) > 1/n\},$$

so

$$A := \bigcup_{n=1}^{\infty} A_n = \{\omega : X_s - \mathbb{E}(X_t | \mathcal{F}_s) > 0\}.$$

Consider  $\mathbb{P}(A) = \mathbb{P}(\bigcup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} \mathbb{P}(A_n)$ . Suppose for some  $n$ ,  $\mathbb{P}(A_n) > \epsilon$ , then note that

$$\begin{aligned} \omega \in A_n & : X_s - \mathbb{E}(X_t | \mathcal{F}_s) > 1/n \\ \omega \in \Omega/A_n & : X_s - \mathbb{E}(X_t | \mathcal{F}_s) \geq 0 \end{aligned}$$

Hence

$$X_s - \mathbb{E}(X_t|\mathcal{F}_s) \geq \frac{1}{n}1_{A_n},$$

taking expectations yields

$$\mathbb{E}(X_s) - \mathbb{E}(X_t) > \frac{\epsilon}{n},$$

but by the constant mean property the left hand side is zero; hence a contradiction, thus all the  $\mathbb{P}(A_n)$  are zero, so

$$X_s = \mathbb{E}(X_t|\mathcal{F}_s) \quad \text{a.s.}$$

□

## 7. The Integral

---

We would like eventually to extend the definition of the integral to integrands which are *previsible* processes and integrators which are *semimartingales* (to be defined later in **these** notes). In fact in these notes we'll only get as far as *continuous semimartingales*; but it is possible to go the whole way and define the integral of a previsible process with respect to a general semimartingale; but some extra problems are thrown up on the way, in particular as regards the construction of the quadratic variation process of a discontinuous process.

Various special classes of process will be needed in the sequel and these are all defined here for convenience. Naturally with terms like 'elementary' and 'simple' occurring many books have different names for the same concepts – so beware!

### 7.1. Elementary Processes

An elementary process  $H_t(\omega)$  is one of the form

$$H_t(\omega) = Z(\omega)1_{(S(\omega), T(\omega)]}(t),$$

where  $S, T$  are stopping times,  $S \leq T \leq \infty$ , and  $Z$  is a bounded  $\mathcal{F}_S$  measurable random variable.

Such a process is the simplest non-trivial example of a *previsible* process. Let's prove that it is previsible:

$H$  is clearly a left continuous process, so we need only show that it is adapted. It can be considered as the pointwise limit of a sequence of right continuous processes

$$H_n(t) = \lim_{n \rightarrow \infty} Z1_{[S_n, T_n)}, \quad S_n = S + \frac{1}{n}, \quad T_n = T + \frac{1}{n}.$$

So it is sufficient to show that  $Z1_{[U, V)}$  is adapted when  $U$  and  $V$  are stopping times which satisfy  $U \leq V$ , and  $Z$  is a bounded  $\mathcal{F}_U$  measurable function. Let  $B$  be a borel set of  $\mathbb{R}$ , then the event

$$\{Z1_{[U, V)}(t) \in B\} = [\{Z \in B\} \cap \{U \leq t\}] \cap \{V > t\}.$$

By the definition of  $U$  as a stopping time and hence the definition of  $\mathcal{F}_U$ , the event enclosed by square brackets is in  $\mathcal{F}_t$ , and since  $V$  is a stopping time  $\{V > t\} = \Omega / \{V \leq t\}$  is also in  $\mathcal{F}_t$ ; hence  $Z1_{[U, V)}$  is adapted.  $\square$

### 7.2. Strictly Simple and Simple Processes

A process  $H$  is *strictly simple* ( $H \in \mathcal{L}^*$ ) if there exist  $0 \leq t_0 \leq \dots \leq t_n < \infty$  and uniformly bounded  $\mathcal{F}_{t_k}$  measurable random variables  $Z_k$  such that

$$H = H_0(\omega)1_0(t) \sum_{k=0}^{n-1} Z_k(\omega)1_{(t_k, t_{k+1}]}(t).$$

[13]

This can be extended to  $H$  is a *simple processes* ( $H \in \mathcal{L}$ ), if there exists a sequence of stopping times  $0 \leq T_0 \leq \dots \leq T_k \rightarrow \infty$ , and  $Z_k$  uniformly bounded  $\mathcal{F}_{T_k}$  measurable random variables such that

$$H = H_0(\omega)1_0(t) + \sum_{k=0}^{\infty} Z_k 1_{(T_k, T_{k+1}]}$$

Similarly a simple process is also a previsible process. The fundamental result will follow from the fact that the  $\sigma$ -algebra generated by the simple processes is exactly the previsible  $\sigma$ -algebra. We shall see the application of this after the next section.

## 8. The Stochastic Integral

---

As has been hinted at earlier the stochastic integral must be built up in stages, and to start with we shall consider integrators which are  $L^2$  bounded martingales, and integrands which are simple processes.

### 8.1. Integral for $H \in \mathcal{L}$ and $M \in \mathcal{M}_2$

For a simple process  $H \in \mathcal{L}$ , and  $M$  an  $L^2$  bounded martingale then the integral may be defined by the ‘martingale transform’ (c.f. discrete martingale theory)

$$\int_0^t H_s dM_s = (H \cdot M)_t := \sum_{k=0}^{\infty} Z_k (M_{T_{k+1} \wedge t} - M_{T_k \wedge t})$$

#### Proposition 8.1.

If  $H$  is a simple function,  $M$  a  $L_2$  bounded martingale, and  $T$  a stopping time. Then

- (i)  $(H \cdot M)^T = (H1_{(0,T]}) \cdot M = H \cdot (M^T)$ .
- (ii)  $(H \cdot M) \in \mathcal{M}_2$ .
- (iii)  $E[(H \cdot M)_\infty^2] = \sum_{k=0}^{\infty} [Z_k^2 (M_{T_{k+1}}^2 - M_{T_k}^2)] \leq \|H\|_\infty^2 E(M_\infty^2)$ .

*Proof*

#### Part (i)

As  $H \in \mathcal{L}$  we can write

$$H = \sum_{k=0}^{\infty} Z_k 1_{(T_k, T_{k+1}]},$$

for  $T_k$  stopping times, and  $Z_k$  an  $\mathcal{F}_{T_k}$  measurable bounded random variable. By our definition for  $M \in \mathcal{M}^2$ , we have

$$(H \cdot M)_t = \sum_{k=0}^{\infty} Z_k (M_{T_{k+1} \wedge t} - M_{T_k \wedge t}),$$

and so, for  $T$  a general stopping time consider  $(H \cdot M)_t^T = (H \cdot M)_{T \wedge t}$  and so

$$(H \cdot M)_t^T = \sum_{k=0}^{\infty} Z_k (M_{T_{k+1} \wedge T \wedge t} - M_{T_k \wedge T \wedge t}).$$

Similar computations can be performed for  $(H \cdot M^T)$ , noting that  $M_t^T = M_{T \wedge t}$  and for  $(H1_{(0,T]} \cdot M)$  yielding the same result in both cases. Hence

$$(H \cdot M)^T = (H1_{(0,T]} \cdot M) = (H \cdot M^T).$$

**Part (ii)**

To prove this result, first we shall establish it for an elementary function  $H \in \mathcal{E}$ , and then extend to  $\mathcal{L}$  by linearity. Suppose

$$H = Z1_{(R,S]},$$

where  $R$  and  $S$  are stopping times and  $Z$  is a bounded  $\mathcal{F}_S$  measurable random variable. Let  $T$  be an arbitrary stopping time. We shall prove that

$$\mathbb{E}((H \cdot M)_T) = \mathbb{E}((H \cdot M)_0),$$

and hence via optional stopping conclude that  $(H \cdot M)_t$  is a martingale.

Note that

$$(H \cdot M)_\infty = Z(M_S - M_R),$$

and hence as  $M$  is a martingale, and  $Z$  is  $\mathcal{F}_R$  measurable we obtain

$$\begin{aligned} \mathbb{E}(H \cdot M)_\infty &= \mathbb{E}(\mathbb{E}(Z(M_S - M_R)) | \mathcal{F}_R) = \mathbb{E}(Z\mathbb{E}((M_S - M_R) | \mathcal{F}_R)) \\ &= 0. \end{aligned}$$

Via part (i) note that  $\mathbb{E}(H \cdot M)_T = \mathbb{E}(H \cdot M^T)$ , so

$$\mathbb{E}(H \cdot M)_T = \mathbb{E}(H \cdot M^T)_\infty = 0.$$

Thus  $(H \cdot M)_t$  is a martingale by optional stopping theorem. By linearity, this result extends to  $H$  a simple function (i.e.  $H \in \mathcal{L}$ ).

**Part (iii)**

We wish to prove that  $(H \cdot M)$  is an  $L^2$  bounded martingale. We again start by considering  $H \in \mathcal{E}$ , an elementary function, i.e.

$$H = Z1_{(R,S]},$$

where as before  $R$  and  $S$  are stopping times, and  $Z$  is a bounded  $\mathcal{F}_R$  measurable random variable.

$$\begin{aligned} \mathbb{E}((H \cdot M)_\infty^2) &= \mathbb{E}(Z^2(M_S - M_R)^2), \\ &= \mathbb{E}(Z^2\mathbb{E}((M_S - M_R)^2 | \mathcal{F}_R)), \end{aligned}$$

where  $Z^2$  is removed from the conditional expectation since it is  $\mathcal{F}_R$  measurable random variable. Using the same argument as used in the orthogonality of martingale increments proof,

$$\mathbb{E}((H \cdot M)_\infty^2) = \mathbb{E}(Z^2\mathbb{E}((M_S^2 - M_R^2) | \mathcal{F}_R)) = \mathbb{E}[(Z^2(M_S^2 - M_R^2))].$$

As  $M$  is an  $L^2$  bounded martingale and  $Z$  is a bounded process,

$$\mathbb{E}((H \cdot M)_\infty^2) \leq \sup_{\omega \in \Omega} 2|Z(\omega)|^2 \mathbb{E}(M_\infty^2).$$

so  $(H \cdot M)$  is an  $L^2$  bounded martingale; so together with part (ii),  $(H \cdot M) \in \mathcal{M}_2$ .

To extend this to simple functions is similar, but requires a little care. In general the orthogonality of increments argument extends to the case where only finitely many of the  $Z_k$  in the definition of the simple function  $H$  are non zero. Let  $K$  be the largest  $k$  such that  $Z_k \neq 0$ .

$$\mathbb{E} \left( (H \cdot M)_\infty^2 \right) = \sum_{k=0}^K \mathbb{E} \left( Z_k^2 \left( M_{T_{k+1}}^2 - M_{T_k}^2 \right) \right),$$

which can be bounded as

$$\begin{aligned} \mathbb{E} \left( (H \cdot M)_\infty^2 \right) &\leq \|H_\infty\|^2 \mathbb{E} \left( \sum_{k=0}^K \left( M_{T_{k+1}}^2 - M_{T_k}^2 \right) \right) \\ &\leq \|H_\infty\|^2 \mathbb{E} \left( M_{T_{K+1}}^2 - M_{T_0}^2 \right) \leq \|H_\infty\|^2 \mathbb{E} M_\infty^2, \end{aligned}$$

since we require  $T_0 = 0$ , and  $M \in \mathcal{M}_2$ , so the final bound is obtained via the  $L^2$  martingale convergence theorem.

Now extend this to the case of an infinite sum; let  $n \leq m$ , we have that

$$(H \cdot M)_{T_m} - (H \cdot M)_{T_n} = (H 1_{(T_n, T_m]} \cdot M),$$

applying the result just proven for finite sums to the right hand side yields

$$\begin{aligned} \|(H \cdot M)_{T_m} - (H \cdot M)_{T_n}\|_2^2 &= \sum_{k=n}^{m-1} \mathbb{E} \left( Z_k^2 \left( M_{T_{k+1}}^2 - M_{T_k}^2 \right) \right) \\ &\leq \|H_\infty\|_2^2 \mathbb{E} \left( M_\infty^2 - M_{T_n}^2 \right). \end{aligned}$$

But by the  $L^2$  martingale convergence theorem the right hand side of this bound tends to zero as  $n \rightarrow \infty$ ; hence  $(H \cdot M)_{T_n}$  converges in  $\mathcal{M}_2$  and the limit must be the pointwise limit  $(H \cdot M)$ . Let  $n = 0$  and  $m \rightarrow \infty$  and the result of part (iii) is obtained.  $\square$

## 8.2. Quadratic Variation

We mentioned earlier that the total variation is the variation which is used by the usual Lebesgue-Stieltjes integral, and that this cannot be used for defining a stochastic integral, since any continuous local martingale of finite variation is indistinguishable from zero. We are now going to look at a variation which will prove fundamental for the construction of the integral. All the definitions as given here aren't based on the partition construction. This is because I shall follow Dellacherie and Meyer and show that the other definitions are equivalent by using the stochastic integral.

### Theorem 8.2.

The quadratic variation process  $\langle M \rangle_t$  of a **continuous**  $L^2$  integrable martingale  $M$  is the unique process  $A_t$  starting from zero such that  $M_t^2 - A_t$  is a uniformly integrable martingale.

*Proof*

For each  $n$  define stopping times

$$S_0^n = 0, \quad S_{k+1}^n = \inf \left\{ t > T_k^n : \left| M_t - M_{T_k^n} \right| > 2^{-n} \right\} \text{ for } k \geq 0$$

Define

$$T_k^n := S_k^n \wedge t$$

Then

$$\begin{aligned} M_t^2 &= \sum_{k \geq 1} \left( M_{t \wedge S_k^n}^2 - M_{t \wedge S_{k-1}^n}^2 \right) \\ &= \sum_{k \geq 1} \left( M_{T_k^n}^2 - M_{T_{k-1}^n}^2 \right) \\ &= 2 \sum_{k \geq 1} M_{T_{k-1}^n} \left( M_{T_k^n} - M_{T_{k-1}^n} \right) + \sum_{k \geq 1} \left( M_{T_k^n} - M_{T_{k-1}^n} \right)^2 \end{aligned} \quad (*)$$

Now define  $H^n$  to be the simple process given by

$$H^n := \sum_{k \geq 1} M_{S_{k-1}^n} 1_{(S_{k-1}^n, S_k^n]}.$$

We can then think of the first term in the decomposition (\*) as  $(H^n \cdot M)$ . Now define

$$A_t^n := \sum_{k \geq 1} \left( M_{T_k^n} - M_{T_{k-1}^n} \right)^2,$$

so the expression (\*) becomes

$$M_t^2 = 2(H^n \cdot M)_t + A_t^n. \quad (**)$$

Note from the construction of the stopping times  $S_k^n$  we have the following properties

$$\begin{aligned} \|H^n - H^{n+1}\|_\infty &= \sup_t |H_t^n - H_t^{n+1}| \leq 2^{-(n+1)} \\ \|H^n - H^{n+m}\|_\infty &= \sup_t |H_t^n - H_t^{n+m}| \leq 2^{-(n+1)} \text{ for all } m \geq 1 \\ \|H^n - M\|_\infty &= \sup_t |H_t^n - M_t| \leq 2^{-n} \end{aligned}$$

Let  $J_n(\omega)$  be the set of all stopping times  $S_k^n(\omega)$  i.e.

$$J_n(\omega) := \{S_k^n(\omega) : k \geq 0\}.$$

Clearly  $J_n(\omega) \subset J_{n+1}(\omega)$ . Now for any  $m \geq 1$ , using proposition 7.1(iii) the following result holds

$$\begin{aligned} \mathbb{E} \left( \left[ (H^n \cdot M) - (H^{n+m} \cdot M) \right]_\infty^2 \right) &= \mathbb{E} \left( \left[ (\{H^n - H^{n+m}\} \cdot M) \right]_\infty^2 \right) \\ &\leq \|H^n - H^{n+m}\|_\infty^2 \mathbb{E}(M_\infty^2) \\ &\leq \left( 2^{-(n+1)} \right)^2 \mathbb{E}(M_\infty^2). \end{aligned}$$

Thus  $(H^n \cdot M)_\infty$  is a Cauchy sequence in the complete Hilbert space  $L^2(\mathcal{F}_\infty)$ ; hence by completeness of the Hilbert Space it converges to a limit in the same space. As  $(H^n \cdot M)$  is a continuous martingale for each  $n$ , so is the limit  $N$  say. By Doob's  $L^2$  inequality applied to the continuous martingale  $(H^n \cdot M) - N$ ,

$$\mathbb{E} \left( \sup_{t \geq 0} |(H^n \cdot M) - N|^2 \right) \leq 4\mathbb{E} \left( [(H \cdot M) - N]_\infty^2 \right) \rightarrow_{n \rightarrow \infty} 0,$$

Hence  $(H^n \cdot M)$  converges to  $N$  uniformly a.s.. From the relation (\*\*) we see that as a consequence of this, the process  $A^n$  converges to a process  $A$ , where

$$M_t^2 = 2N_t + A_t.$$

Now we must check that this limit process  $A$  is increasing. Clearly  $A^n(S_k^n) \leq A^n(S_{k+1}^n)$ , and since  $J_n(\omega) \subset J_{n+1}(\omega)$ , it is also true that  $A(S_k^n) \leq A(S_{k+1}^n)$  for all  $n$  and  $k$ , and so  $A$  is certainly increasing on the closure of  $J(\omega) := \cup_n J_n(\omega)$ . However if  $I$  is an open interval in the complement of  $J$ , then no stopping time  $S_k^n$  lies in this interval, so  $M$  must be constant throughout  $I$ , so the same is true for the process  $A$ . Hence the process  $A$  is continuous, increasing, and null at zero; such that  $M_t^2 - A_t = 2N_t$ , where  $N_t$  is a UI martingale (since it is  $L^2$  bounded). Thus we have established the existence result. It only remains to consider uniqueness.

Uniqueness follows from the result that a continuous local martingale of finite variation is everywhere zero. Suppose the process  $A$  in the above definition were not unique. That is suppose that also for some  $B_t$  continuous increasing from zero,  $M_t^2 - B_t$  is a UI martingale. Then as  $M_t^2 - A_t$  is also a UI martingale by subtracting these two equations we get that  $A_t - B_t$  is a UI martingale, null at zero. It clearly must have finite variation, and hence be zero.  $\square$

The following corollary will be needed to prove the integration by parts formula, and can be skipped on a first reading; however it is clearer to place it here, since this avoids having to redefine the notation.

**Corollary 8.3.**

Let  $M$  be a bounded continuous martingale, starting from zero. Then

$$M_t^2 = 2 \int_0^t M_s dM_s + \langle M \rangle_t.$$

*Proof*

In the construction of the quadratic variation process the quadratic variation was constructed as the uniform limit in  $L^2$  of processes  $A_t^n$  such that

$$A_t^n = M_t^2 - 2(H^n \cdot M)_t,$$

where each  $H^n$  was a bounded previsible process, such that

$$\sup_t |H_t^n - M| \leq 2^{-n},$$

and hence  $H^n \rightarrow M$  in  $L^2(M)$ , so the martingales  $(H^n \cdot M)$  converge to  $(M \cdot M)$  uniformly in  $L^2$ , hence it follows immediately that

$$M_t^2 = 2 \int_0^t M_s dM_s + \langle M \rangle_t,$$

□

**Theorem 8.4.**

The quadratic variation process  $\langle M \rangle_t$  of a **continuous** local martingale  $M$  is the unique increasing process  $A$ , starting from zero such that  $M^2 - A$  is a local martingale.

*Proof*

We shall use a localisation technique to extend the definition of quadratic variation from  $L^2$  bounded martingales to general local martingales.

The mysterious seeming technique of localisation isn't really that complex to understand. The idea is that it enables us to extend a definition which applies for 'X widgets' to one valid for 'local X widgets'. It achieves this by using a sequence of stopping times which reduce the 'local X widgets' to 'X widgets'; the original definition can then be applied to the stopped version of the 'X widget'. We only need to check that we can sew up the pieces without any holes i.e. that our definition is independent of the choice of stopping times!

Let  $T_n = \inf\{t : |M_t| > n\}$ , define a sequence of stopping times. Now define

$$\langle M \rangle_t := \langle M^{T_n} \rangle \text{ for } 0 \leq t \leq T_n$$

To check the consistency of this definition note that

$$\langle M^{T_n} \rangle_{T_{n-1}} = \langle M^{T_{n-1}} \rangle$$

and since the sequence of stopping times  $T_n \rightarrow \infty$ , we see that  $\langle M \rangle$  is defined for all  $t$ . Uniqueness follows from the result that any finite variation continuous local martingale starting from zero is identically zero. □

The quadratic variation turns out to be the 'right' sort of variation to consider for a martingale; since we have already shown that all but the zero martingale have infinite total variation; and it can be shown that the higher order variations of a martingale are zero a.s.. Note that the definition given is for a **continuous local martingale**; we shall see later how to extend this to a continuous semimartingale.

### 8.3. Covariation

From the definition of the quadratic variation of a local martingale we can define the covariation of two local martingales  $N$  and  $M$  which are locally  $L^2$  bounded via the *polarisation identity*

$$\langle M, N \rangle := \frac{\langle M + N \rangle - \langle M - N \rangle}{4}.$$

We need to generalise this slightly, since the above definition required the quadratic variation terms to be finite. We can prove the following theorem in a straightforward

manner using the definition of quadratic variation above, and this will motivate the general definition of the *covariation* process.

**Theorem 8.5.**

For  $M$  and  $N$  two local martingales which are locally  $L^2$  bounded then there exists a unique finite variation process  $A$  starting from zero such that  $MN - A$  is a local martingale. This process  $A$  is the covariation of  $M$  and  $N$ .

This theorem is turned round to give the usual definition of the covariation process of two continuous local martingales as:

**Definition 8.6.**

For two continuous local martingales  $N$  and  $M$ , there exists a unique finite variation process  $A$ , such that  $MN - A$  is a local martingale. The covariance process of  $N$  and  $M$  is defined as this process  $A$ .

It can readily be verified that the covariation process can be regarded as a symmetric bilinear form on the space of local martingales, i.e. for  $L, M$  and  $N$  continuous local martingales

$$\begin{aligned}\langle M + N, L \rangle &= \langle M, L \rangle + \langle N, L \rangle, \\ \langle M, N \rangle &= \langle N, M \rangle, \\ \langle \lambda M, N \rangle &= \lambda \langle M, N \rangle, \quad \lambda \in \mathbb{R}.\end{aligned}$$

#### 8.4. Extension of the Integral to $L^2(M)$

We have previously defined the integral for  $H$  a simple function (in  $\mathcal{L}$ ), and  $M \in \mathcal{M}_2^c$ , and we have noted that  $(H \cdot M)$  is itself in  $\mathcal{M}_2$ . Hence

$$\mathbb{E}((H \cdot M)_\infty^2) = \mathbb{E}\left(Z_{i-1}^2 (M_{T_i} - M_{T_{i-1}})^2\right)$$

Recall that for  $M \in \mathcal{M}_2$ , then  $M^2 - \langle M \rangle$  is a uniformly integrable martingale. Hence for  $S$  and  $T$  stopping times such that  $S \leq T$ , then

$$\mathbb{E}((M_T - M_S)^2 | \mathcal{F}_S) = \mathbb{E}(M_T^2 - M_S^2 | \mathcal{F}_S) = \mathbb{E}(\langle M \rangle_T - \langle M \rangle_S | \mathcal{F}_S).$$

So summing we obtain

$$\begin{aligned}\mathbb{E}((H \cdot M)_\infty^2) &= \mathbb{E} \sum Z_{i-1}^2 (\langle M \rangle_{T_i} - \langle M \rangle_{T_{i-1}}), \\ &= \mathbb{E}((H^2 \cdot \langle M \rangle)_\infty).\end{aligned}$$

In the light of this, we define a seminorm  $\|H\|_M$  via

$$\|H\|_M = \left[ \mathbb{E}((H^2 \cdot \langle M \rangle)_\infty) \right]^{1/2} = \left[ \mathbb{E} \left( \int_0^\infty H_s^2 d\langle M \rangle_s \right) \right]^{1/2}.$$

The space  $\mathcal{L}^2(M)$  is then defined as the subspace of the previsible processes, where this seminorm is finite, i.e.

$$\mathcal{L}^2(M) := \{\text{previsible processes } H \text{ such that } \|H\|_M < \infty\}.$$

However we would actually like to be able to treat this as a Hilbert space, and there remains a problem, namely that if  $X \in \mathcal{L}^2(M)$  and  $\|X\|_M = 0$ , this doesn't imply that  $X$  is the zero process. Thus we follow the usual route of defining an equivalence relation via  $X \sim Y$  if and only if  $\|X - Y\|_M = 0$ . We now define

$$L^2(M) := \{\text{equivalence classes of previsible processes } H \text{ such that } \|H\|_M < \infty\},$$

and this is a Hilbert space with norm  $\|\cdot\|_M$  (it can be seen that it is a Hilbert space by considering it as suitable  $L^2$  space).

This establishes an isometry (called the *Itô isometry*) between the spaces  $L^2(M) \cap \mathcal{L}$  and  $L^2(\mathcal{F}_\infty)$  given by

$$\begin{aligned} I : L^2(M) \cap \mathcal{L} &\rightarrow L^2(\mathcal{F}_\infty) \\ I : H &\mapsto (H \cdot M)_\infty \end{aligned}$$

Remember that there is a basic bijection between the space  $\mathcal{M}_2$  and the Hilbert Space  $L^2(\mathcal{F}_\infty)$  in which each square integrable martingale  $M$  is represented by its limiting value  $M_\infty$ , so the image under the isometry  $(H \cdot M)_\infty$  in  $L^2(\mathcal{F}_\infty)$  may be thought of as describing an  $\mathcal{M}_2$  martingale. Hence this endows  $\mathcal{M}_2$  with a Hilbert Space structure, with an inner product given by

$$(M, N) = \mathbb{E}(N_\infty M_\infty).$$

We shall now use this Itô isometry to extend the definition of the stochastic integral from  $\mathcal{L}$  (the class of simple functions) to the whole of  $L^2(M)$ . Roughly speaking we shall approximate an element of  $L^2(M)$  via a sequence of simple functions converging to it; just as in the construction of the Lebesgue Integral. In doing this, we shall use the *Monotone Class Theorem*.

Recall that in the conventional construction of the Lebesgue integration, and proof of the elementary results the following standard machine is repeatedly invoked. To prove a 'linear' result for all  $h \in L^1(S, \Sigma, \mu)$ , proceed in the following way:

- (i) Show the result is true for  $h$  and indicator function.
- (ii) Show that by linearity the result extends to all positive step functions.
- (iii) Use the Monotone convergence theorem to see that if  $h_n \uparrow h$ , where the  $h_n$  are step functions, then the result must also be true for  $h$  a non-negative,  $\Sigma$  measurable function.
- (iv) Write  $h = h^+ - h^-$  where both  $h^+$  and  $h^-$  are non-negative functions and use linearity to obtain the result for  $h \in L^1$ .

The monotone class lemmas is a replacement for this procedure, which hides away all the 'machinery' used in the constructions.

### Monotone Class Theorem.

Let  $\mathcal{A}$  be  $\pi$ -system generating the  $\sigma$ -algebra  $\mathcal{F}$  (i.e.  $\sigma(\mathcal{A}) = \mathcal{F}$ ). If  $\mathcal{H}$  is a linear set of bounded functions from  $\Omega$  to  $\mathbb{R}$  satisfying

- (i)  $1_A \in \mathcal{H}$ , for all  $A \in \mathcal{A}$ ,

(ii)  $0 \leq f_n \uparrow f$ , where  $f_n \in \mathcal{H}$  and  $f$  is a bounded function  $f : \Omega \rightarrow \mathbb{R}$ , then this implies that  $f \in \mathcal{H}$ ,

then  $\mathcal{H}$  contains every bounded,  $\mathcal{F}$ -measurable function  $f : \Omega \rightarrow \mathbb{R}$ .

In order to apply this in our case, we need to prove that the  $\sigma$ -algebra of previsible processes is that generated by the simple functions.

### The Previsible $\sigma$ -field and the Simple Processes

It is fairly simple to show that the space of simple processes  $\mathcal{L}$  forms a vector space (exercise: check linearity, constant multiples and zero).

#### Lemma 8.7.

The  $\sigma$ -algebra generated by the simple functions is the previsible  $\sigma$ -algebra i.e. the previsible  $\sigma$ -algebra is the smallest  $\sigma$ -algebra with respect to which every simple process is measurable.

*Proof*

It suffices to show that every left continuous right limit process, which is bounded and adapted to  $\mathcal{F}_t$  is measurable with respect to the  $\sigma$ -algebra generated by the simple functions. Let  $H_t$  be a bounded left continuous right limits process, then

$$H = \lim_{k \rightarrow \infty} \lim_{n \rightarrow \infty} \sum_{i=2}^{nk} H_{(i-1)/n} \left( \frac{i-1}{n}, \frac{i}{n} \right],$$

and if  $H_t$  is adapted to  $\mathcal{F}_t$  then  $H_{(i-1)/n}$  is a bounded element of  $\mathcal{F}_{(i-1)/n}$ .

We can now apply the Monotone Class Theorem to the vector space  $\mathcal{H}$  of processes with a time parameter in  $(0, \infty)$ , regarded as maps from  $(0, \infty) \times \Omega \rightarrow \mathbb{R}$ . Then if this vector space contains all the simple functions i.e.  $\mathcal{L} \subset \mathcal{H}$ , then  $\mathcal{H}$  contains every bounded previsible process on  $(0, \infty)$ .

### Assembling the Pieces

Since  $I$  is an isometry it has a unique extension to the closure of

$$\mathcal{U} = L^2(M) \cap \mathcal{L},$$

in  $L^2(M)$ . By the application of the monotone class lemma to  $\mathcal{H} = \overline{\mathcal{U}}$ , and the  $\pi$ -system of simple functions. We see that  $\overline{\mathcal{U}}$  must contain every bounded previsible process; hence  $\overline{\mathcal{U}} = L^2(M)$ . Thus the Itô Isometry extends to a map from  $L^2(M)$  to  $L^2(\mathcal{F}_\infty)$ .

Let us look at this result more informally. For a previsible  $H \in L^2(M)$ , because of the density of  $\mathcal{L}$  in  $L^2(M)$ , we can find a sequence of simple functions  $H_n$  which converges to  $H$ , as  $n \rightarrow \infty$ . We then consider  $I(H)$  as the limit of the  $I(H_n)$ . To verify that this limit is unique, suppose that  $H'_n \rightarrow H$  as  $n \rightarrow \infty$  also, where  $H'_n \in \mathcal{L}$ . Note that  $H_n - H'_n \in \mathcal{L}$ . Also  $H_n - H'_n \rightarrow 0$  and so  $((H_n - H'_n) \cdot M) \rightarrow 0$ , and hence by the Itô isometry the limits  $\lim_{n \rightarrow \infty} (H_n \cdot M)$  and  $\lim_{n \rightarrow \infty} (H'_n \cdot M)$  coincide.

The following result is essential in order to extend the integral to continuous local martingales.

**Proposition 8.8.**

For  $M \in \mathcal{M}_2$ , for any  $H \in L^2(M)$  and for any stopping time  $T$  then

$$(H \cdot M)^T = (H1_{(0,T]} \cdot M) = (H \cdot M^T).$$

*Proof*

Consider the following linear maps in turn

$$\begin{aligned} f_1 : L^2(\mathcal{F}_\infty) &\rightarrow L^2(\mathcal{F}_\infty) \\ f_1 : Y &\mapsto \mathbb{E}(Y|\mathcal{F}_T) \end{aligned}$$

This map is a contraction on  $L^2(\mathcal{F}_\infty)$  since by conditional Jensen's inequality

$$\mathbb{E}(Y_\infty|\mathcal{F}_T)^2 \leq \mathbb{E}(Y_\infty^2|\mathcal{F}_T),$$

and taking expectations yields

$$\|\mathbb{E}(Y|\mathcal{F}_T)\|_2^2 = \mathbb{E}(\mathbb{E}(Y_\infty|\mathcal{F}_T)^2) \leq \mathbb{E}(\mathbb{E}(Y_\infty^2|\mathcal{F}_T)) = \mathbb{E}(Y_\infty^2) = \|Y\|_2^2.$$

Hence  $f_1$  is a contraction on  $L^2(\mathcal{F}_\infty)$ . Now

$$\begin{aligned} f_2 : L^2(M) &\rightarrow L^2(M) \\ f_2 : H &\mapsto H1_{(0,T]} \end{aligned}$$

Clearly from the definition of  $\|\cdot\|_M$ , and from the fact that the quadratic variation process is increasing

$$\|H1_{(0,T]}\|_M = \int_0^\infty H_s^2 1_{(0,T]} d\langle M \rangle_s = \int_0^T H_s^2 d\langle M \rangle_s \leq \int_0^\infty H_s^2 d\langle M \rangle_s = \|H\|_M.$$

Hence  $f_2$  is a contraction on  $L^2(M)$ . Hence if  $I$  denotes the Itô isometry then  $f_1 \circ I$  and  $I \circ f_2$  are also contractions from  $L^2(M)$  to  $L^2(\mathcal{F}_\infty)$ , (using the fact that  $I$  is an isometry between  $L^2(M)$  and  $L^2(\mathcal{F}_\infty)$ ).

Now introduce  $I^{(T)}$ , the stochastic integral map associated with  $M^T$ , i.e.

$$I^{(T)}(H) \equiv (H \cdot M^T)_\infty.$$

Note that

$$\|I^{(T)}(H)\|_2 = \|H\|_{M^T} \leq \|H\|_M.$$

We have previously shown that the maps  $f_1 \circ I$  and  $I \circ f_2$  and  $H \mapsto I^{(T)}(H)$  agree on the space of simple functions by direct calculation. We note that  $\mathcal{L}$  is dense in  $L^2(M)$  (from application of Monotone Class Lemma to the simple functions). Hence from the three bounds above the three maps agree on  $L^2(M)$ .  $\square$

### 8.5. Localisation

We've already met the idea of localisation in extending the definition of quadratic variation from  $L^2$  bounded continuous martingales to continuous local martingales. In this context a previsible process  $\{H_t\}_{t \geq 0}$ , is *locally previsible* if there exists a sequence of stopping times  $T_n \rightarrow \infty$  such that for all  $n$   $H1_{(0, T_n]}$  is a previsible process. Fairly obviously every previsible process has this property. However if in addition we want the process  $H$  to be *locally bounded* we need the condition that there exists a sequence  $T_n$  of stopping times, tending to infinity such that  $H1_{(0, T_n]}$  is uniformly bounded for each  $n$ .

For the integrator (a martingale of integrable variation say), the localisation is to a local martingale, that is one which has a sequence of stopping times  $T_n \rightarrow \infty$  such that for all  $n$ ,  $X^{T_n}$  is a genuine martingale.

If we can prove a result like

$$(H \cdot X)^T = (H1_{(0, T]} \cdot X^T)$$

for  $H$  and  $X$  in their original (i.e. non-localised classes) then it is possible to *extend* the definition of  $(H \cdot X)$  to the local classes.

Note firstly that for  $H$  and  $X$  local, and  $T_n$  a reducing sequence<sup>1</sup> of stopping times for both  $H$  and  $X$  then we see that  $(H1_{(0, T]} \cdot X^T)$  is defined in the existing fashion. Also note that if  $T = T_{n-1}$  we can check consistency

$$(H1_{(0, T_n]} \cdot X^{T_n})^{T_{n-1}} = (H \cdot X)^{T_{n-1}} = (H1_{(0, T_{n-1}]} \cdot X^{T_{n-1}}).$$

Thus it is consistent to define  $(H \cdot X)_t$  on  $t \in [0, \infty)$  via

$$(H \cdot X)^{T_n} = (H1_{(0, T_n]} \cdot X^{T_n}), \quad \forall n.$$

We must check that this is well defined, viz if we choose another regularising sequence  $S_n$ , we get the same definition of  $(H \cdot X)$ . To see this note:

$$(H1_{(0, T_n]} \cdot X^{T_n})^{S_n} = (H1_{(0, T_n \wedge S_n]} \cdot X^{T_n \wedge S_n}) = (H1_{(0, S_n]} \cdot X^{S_n})^{T_n},$$

hence the definition of  $(H \cdot X)_t$  is the same if constructed from the regularising sequence  $S_n$  as if constructed via  $T_n$ .

### 8.6. Some Important Results

We can now extend most of our results to stochastic integrals of a previsible process  $H$  with respect to a **continuous** local martingale  $M$ . In fact in these notes we will never drop the continuity requirement. It can be done; but it requires considerably more work, especially with regard to the definition of the quadratic variation process.

---

<sup>1</sup> The reducing sequence is the sequence of stopping times tending to infinity which makes the local version of the object into the non-local version. We can find one such sequence, because if say  $\{T_n\}$  reduces  $H$  and  $\{S_n\}$  reduces  $X$  then  $T_n \wedge S_n$  reduces both  $H$  and  $X$ .

**Theorem 8.9.**

Let  $H$  be a locally bounded previsible process, and  $M$  a continuous local martingale. Let  $T$  be an arbitrary stopping time. Then:

- (i)  $(H \cdot M)^T = (H1_{(0,T]} \cdot M) = (H \cdot M^T)$
- (ii)  $(H \cdot M)$  is a continuous local martingale
- (iii)  $\langle H \cdot M \rangle = H^2 \cdot \langle M \rangle$
- (iv)  $H \cdot (K \cdot M) = (HK) \cdot M$

*Proof*

The proof of parts (i) and (ii) follows from the result used in the localisation that:

$$(H \cdot M)^T = (H1_{(0,T]} \cdot M) = (H \cdot M^T)$$

for  $H$  bounded previsible process in  $L^2(M)$  and  $M$  an  $L^2$  bounded martingale. Using this result it suffices to prove (iii) and (iv) where  $M$ ,  $H$  and  $K$  are **uniformly** bounded (via localisation).

**Part (iii)**

$$\begin{aligned} \mathbb{E} [(H \cdot M)_T^2] &= \mathbb{E} [(H1_{(0,T]} \cdot M) \cdot M]_\infty^2 \\ &= \mathbb{E} [(H1_{(0,T]} \cdot \langle M \rangle)_\infty^2] \\ &= \mathbb{E} [(H^2 \cdot \langle M \rangle)_T] \end{aligned}$$

Hence we see that  $(H \cdot M)^2 - (H^2 \cdot \langle M \rangle)$  is a martingale (via the optional stopping theorem), and so by uniqueness of the quadratic variation process, we have established

$$\langle H \cdot M \rangle = H^2 \cdot \langle M \rangle.$$

**Part (iv)**

The truth of this statement is readily established for  $H$  and  $K$  simple functions (in  $\mathcal{L}$ ). To extend to  $H$  and  $K$  bounded previsible processes note that

$$\begin{aligned} \mathbb{E} [(H \cdot (K \cdot M))_\infty^2] &= \mathbb{E} [(H^2 \cdot \langle K \cdot M \rangle)_\infty] \\ &= \mathbb{E} [(H^2 \cdot (K^2 \cdot \langle M \rangle))_\infty] \\ &= \mathbb{E} [((HK)^2 \cdot \langle M \rangle)_\infty] \\ &= \mathbb{E} [((HK) \cdot M)_\infty^2] \end{aligned}$$

Also note the following bound

$$\mathbb{E} [((HK)^2 \cdot \langle M \rangle)_\infty] \leq \min \{ \|H\|_\infty^2 \|K\|_M^2, \|H\|_M^2 \|K\|_\infty^2 \}.$$

□

## 9. Semimartingales

---

I mentioned at the start of these notes that the most general form of the stochastic integral would have a previsible process as the integrand and a semimartingale as an integrator. Now it's time to extend the definition of the Itô integral to the case of semimartingale integrators.

**Definition 9.1.**

A process  $X$  is a semimartingale if  $X$  is an adapted CADLAG process which has a decomposition

$$X = X_0 + M + A,$$

where  $M$  is a local martingale, null at zero and  $A$  is a process null at zero, with paths of finite variation.

Note that the decomposition is **not** necessarily unique as there exist martingales which have finite variation. To remove many of these difficulties we shall impose a continuity condition, since under this most of our problems will vanish.

**Definition 9.2.**

A continuous semimartingale is a process  $(X_t)_{t \geq 0}$  which has a Doob-Meyer decomposition

$$X = X_0 + M + A,$$

where  $X_0$  is  $\mathcal{F}_0$ -measurable,  $M_0 = A_0 = 0$ ,  $M_t$  is a continuous local martingale and  $A_t$  is a continuous adapted process of finite variation.

**Theorem 9.3.**

The Doob-Meyer decomposition in the definition of a continuous semimartingale is unique.

*Proof*

Let another such decomposition be

$$X = X_0 + M' + A',$$

where  $M'$  is a continuous local martingale and  $A$  a continuous adapted process of finite variation. Then consider the process  $N$ , where

$$N = M' - M = A' - A,$$

by the first equality,  $N$  is the difference of two continuous local martingales, and hence is itself a continuous local martingale; and by the second inequality it has finite variation. Hence by an earlier proposition (5.2) it must be zero. Hence  $M' = M$  and  $A' = A$ .  $\square$

We **define**† the quadratic variation of the continuous semimartingale as that of the continuous local martingale part i.e. for  $X = X_0 + M + A$ ,

$$\langle X \rangle := \langle M \rangle.$$

---

† These definitions can be made to look natural by considering the quadratic variation defined in terms of a sum of squared increments; but following this approach, these are result which are proved later using the Itô integral, since this provided a better approach to the discontinuous theory.

Similarly if  $Y + Y_0 + N + B$  is another semimartingale, where  $B$  is finite variation and  $N$  is a continuous local martingale, we define

$$\langle X, Y \rangle := \langle M, N \rangle.$$

We can extend the definition of the stochastic integral to continuous semimartingale integrators by defining

$$(H \cdot X) := (H \cdot M) + (H \cdot A),$$

where the first integral is a stochastic integral as defined earlier and the second is a Lebesgue-Stieltjes integral (as the integrator is a process of finite variation).

## 10. Relations to Sums

---

This section is optional; and is included to bring together the two approaches to the constructions involved in the stochastic integral.

For example the quadratic variation of a process can either be defined in terms of martingale properties, or alternatively in terms of sums of squares of increments.

### 10.1. The UCP topology

We shall meet the notion of convergence uniformly on compacts in probability when considering stochastic integrals as limits of sums, so it makes sense to review this topology here.

**Definition 10.1.**

A sequence  $\{H_n\}_{n \geq 1}$  converges to a process  $H$  uniformly on compacts in probability (abbreviated u.c.p.) if for each  $t > 0$ ,

$$\sup_{0 \leq s \leq t} |H_s^n - H_s| \rightarrow 0 \text{ in probability.}$$

At first sight this may seem to be quite an esoteric definition; in fact it is a natural extension of convergence in probability to processes. It would also appear to be quite difficult to handle, however Doob's martingale inequalities provide the key to handling it. Let

$$H_t^* = \sup_{0 \leq s \leq t} |H_s|,$$

then for  $Y^n$  a CADLAG process,  $Y^n$  converges to  $Y$  u.c.p. iff  $(Y^n - Y)^*$  converges to zero in probability for each  $t \geq 0$ . Thus to prove that a sequence converges u.c.p. it often suffices to apply Doob's inequality to prove that the supremum converges to zero in  $L^2$ , whence it must converge to zero in probability, whence u.c.p. convergence follows.

The space of CADLAG processes with u.c.p. topology is in fact metrizable, a compatible metric is given by

$$d(X, Y) = \sum_{n=1}^{\infty} \frac{1}{2^n} \mathbb{E}(\min(1, (X - Y)_n^*)),$$

for  $X$  and  $Y$  CADLAG processes. The metric space can also be shown to be complete. For details see Protter.

Since we have just met a new kind of convergence, it is helpful to recall the other usual types of convergence on a probability space. For convenience here are the usual definitions:

**Pointwise**

A sequence of random variables  $X_n$  converges to  $X$  pointwise if for all  $\omega$  not in some null set,

$$X_n(\omega) \rightarrow X(\omega).$$

**Probability**

A sequence of r.v.s  $X_n$  converges to  $X$  in probability, if for any  $\epsilon > 0$ ,

$$\mathbb{P}(|X_n - X| > \epsilon) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

**$L^p$  convergence**

A sequence of random variables  $X_n$  converges to  $X$  in  $L^p$ , if

$$\mathbb{E}|X_n - X|^p \rightarrow 0, \text{ as } n \rightarrow \infty.$$

It is trivial to see that pointwise convergence implies convergence in probability. It is also true that  $L^p$  convergence implies convergence in probability as the following theorem shows

**Theorem 10.2.**

If  $X_n$  converges to  $X$  in  $L^p$  for  $p > 0$ , then  $X_n$  converges to  $X$  in probability.

*Proof*

Apply Chebyshev's inequality to  $f(x) = x^p$ , which yields for any  $\epsilon > 0$ ,

$$\mathbb{P}(|X_n| \geq \epsilon) \leq \epsilon^{-p} \mathbb{E}(|X_n|^p) \rightarrow 0, \text{ as } n \rightarrow \infty.$$

□

**Theorem 10.3.**

If  $X_n \rightarrow X$  in probability, then there exists a subsequence  $n_k$  such that  $X_{n_k} \rightarrow X$  a.s.

**Theorem 10.4.**

If  $X_n \rightarrow X$  a.s., then  $X_n \rightarrow X$  in probability.

## 10.2. Approximation via Riemann Sums

Following Dellacherie and Meyer we shall establish the equivalence of the two constructions for the quadratic variation by the following theorem which approximates the stochastic integral via Riemann sums.

**Theorem 10.2.**

Let  $X$  be a semimartingale, and  $H$  a locally bounded previsible CADLAG process starting from zero. Then

$$\int_0^t H_s dX_s = \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} H_{t \wedge k2^{-n}} (X_{t \wedge (k+1)2^{-n}} - X_{t \wedge k2^{-n}}) \text{ u.c.p.}$$

*Proof*

Let  $K_s = H_s 1_{s \leq t}$ , and define the following sequence of simple function approximations

$$K_s^n := \sum_{k=0}^{\infty} H_{t \wedge k2^{-n}} 1_{(t \wedge k2^{-n}, t \wedge (k+1)2^{-n}]}(s).$$

Clearly this sequence  $K_s^n$  converges pointwise to  $K_s$ . We can decompose the semimartingale  $X$  as  $X = X_0 + A_t + M_t$  where  $A_t$  is of finite variation and  $M_t$  is a continuous local martingale, both starting from zero. The result that

$$\int_0^t K_s^n dA_s \rightarrow \int_0^t K_s dA_s, \text{ u.c.p.}$$

is standard from the Lebesgue-Stieltjes theory. Let  $T_k$  be a reducing sequence for the continuous local martingale  $M$  such that  $M^{T_k}$  is a **bounded** martingale. Also since  $K$  is locally bounded we can find a sequence of stopping times  $S_k$  such that  $K^{S_k}$  is a bounded previsible process. It therefore suffices to prove for a sequence of stopping times  $R_k$  such that  $R_k \uparrow \infty$ , then

$$(K^n \cdot M)_s^{R_k} \rightarrow (K \cdot M)_s^{R_k}, \text{ u.c.p..}$$

By Doob's  $L^2$  inequality, and the Itô isometry we have

$$\begin{aligned} \mathbb{E} [((K^n \cdot M) - (K \cdot M))^*]^2 &\leq 4\mathbb{E} [(K^n \cdot M) - (K \cdot M)]^2, && \text{Doob } L^2 \\ &\leq 4\|K^n - K\|_M^2, && \text{Itô Isometry} \\ &\leq 4 \int (K_s^n - K_s)^2 d\langle M \rangle_s \end{aligned}$$

As  $|K^n - K| \rightarrow 0$  pointwise, and  $K$  is bounded, clearly  $|K^n - K|$  is also bounded uniformly in  $n$ . Hence by the Dominated Convergence Theorem for the Lebesgue-Stieltjes integral

$$\int (K_s^n - K_s)^2 d\langle M \rangle_s \rightarrow 0 \text{ a.s..}$$

Hence, we may conclude

$$\mathbb{E} [((K^n \cdot M) - (K \cdot M))^*]^2 \rightarrow 0, \text{ as } n \rightarrow \infty.$$

So

$$[(K^n \cdot M) - (K \cdot M)]^* \rightarrow 0 \text{ in } L^2,$$

as  $n \rightarrow \infty$ ; but this implies that

$$[(K^n \cdot M) - (K \cdot M)]^* \rightarrow 0 \text{ in probability.}$$

Hence

$$[(K^n \cdot M) - (K \cdot M)] \rightarrow 0 \text{ u.c.p.}$$

as required, and putting the two parts together yields

$$\int_0^t K_s^n dX_s \rightarrow \int_0^t K_s dX_s, \text{ u.c.p.}$$

which is the required result.  $\square$

This result can now be applied to the construction of the quadratic variation process, as illustrated by the next theorem.

**Theorem 10.3.**

The quadratic variation process  $\langle X \rangle_t$  is equal to the following limit in probability

$$\langle X \rangle_t = \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} (X_{t \wedge (k+1)2^{-n}} - X_{t \wedge k2^{-n}})^2 \text{ in probability.}$$

*Proof*

In the theorem (7.2) establishing the existence of the quadratic variation process, we noted in (\*\*) that

$$A_t^n = M_t^2 - 2(H^n \cdot M)_t.$$

Now from application of the previous theorem

$$2 \int_0^t X_s dX_s = \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} X_{t \wedge k2^{-n}} (X_{t \wedge (k+1)2^{-n}} - X_{t \wedge k2^{-n}}).$$

In addition,

$$X_t^2 - X_0^2 = \sum_{k=0}^{\infty} (X_{t \wedge (k+1)2^{-n}}^2 - X_{t \wedge k2^{-n}}^2).$$

The difference of these two equations yields

$$A_t = X_0^2 + \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} (X_{t \wedge (k+1)2^{-n}} - X_{t \wedge k2^{-n}})^2,$$

where the limit is taken in probability. Hence the function  $A$  is increasing and positive on the rational numbers, and hence on the whole of  $\mathbb{R}$  by right continuity.  $\square$

**Remark**

The theorem can be strengthened still further by a result of Doléans-Dade to the effect that for  $X$  a continuous semimartingale

$$\langle X \rangle_t = \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} (X_{t \wedge (k+1)2^{-n}} - X_{t \wedge k2^{-n}})^2,$$

where the limit is in the strong sense in  $L^1$ . This result is harder to prove (essentially the uniform integrability of the sums must be proven) and this is not done here.

## 11. Itô's Formula

---

Itô's Formula is the analog of integration by parts in the stochastic calculus. It is also the first place where we see a major difference creep into the theory, and realise that our formalism has found a new subtlety in the subject.

More importantly, it is the fundamental weapon used to evaluate Itô integrals; we shall see some examples of this shortly.

The Itô isometry provides a clean-cut definition of the stochastic integral; however it was originally defined via the following theorem of Kunita and Watanabe.

**Theorem (Kunita-Watanabe Identity) 11.1.**

Let  $M \in \mathcal{M}_2$  and  $H$  and  $K$  are locally bounded previsible processes. Then  $(H \cdot M)_\infty$  is the unique element of  $L^2(\mathcal{F}_\infty)$  such that for every  $N \in \mathcal{M}_2$  we have:

$$\mathbb{E}[(H \cdot M)_\infty N_\infty] = \mathbb{E}[(H \cdot \langle M, N \rangle)_\infty]$$

Moreover we have

$$\langle (H \cdot M), N \rangle = H \cdot \langle M, N \rangle.$$

*Proof*

Consider an elementary function  $H$ , so  $H = Z1_{(S,T]}$ , where  $Z$  is an  $\mathcal{F}_S$  measurable bounded random variable, and  $S$  and  $T$  are stopping times such that  $S \leq T$ . It is clear that

$$\begin{aligned} \mathbb{E}[(H \cdot M)_\infty N_\infty] &= \mathbb{E}[Z(M_T - M_S)N_\infty] \\ &= \mathbb{E}[Z(M_T N_T - M_S N_S)] \\ &= \mathbb{E}[M_\infty(H \cdot N)_\infty] \end{aligned}$$

Now by linearity this can be extended to establish the result for all simple functions (in  $\mathcal{L}$ ). We finally extend to general locally bounded previsible  $H$ , by considering a sequence (provided it exists) of simple functions  $H^n$  such that  $H^n \rightarrow H$  in  $L^2(M)$ . Then there exists a subsequence  $n_k$  such that  $H^{n_k}$  converges to  $H$  in  $L^2(N)$ . Then

$$\begin{aligned} \mathbb{E}((H^{n_k} \cdot M)_\infty N_\infty) - \mathbb{E}((H \cdot M)_\infty N_\infty) &= \mathbb{E}\left(\left((H^{n_k} - H) \cdot M\right) N_\infty\right) \\ &\leq \sqrt{\mathbb{E}\left([\left(H^{n_k} - H\right) \cdot M\right]^2\right)} \sqrt{\mathbb{E}(N_\infty^2)} \\ &\leq \sqrt{\mathbb{E}\left([\left(H^{n_k} \cdot M\right) - (H \cdot M)]^2\right)} \sqrt{\mathbb{E}(N_\infty^2)} \end{aligned}$$

By construction  $H^{n_k} \rightarrow H$  in  $L^2(M)$  which means that

$$\|H^{n_k} - H\|_M \rightarrow 0, \text{ as } k \rightarrow \infty.$$

By the Itô isometry

$$\mathbb{E}\left[\left((H^{n_k} - H) \cdot M\right)^2\right] = \|H^{n_k} - H\|_M^2 \rightarrow 0, \text{ as } k \rightarrow \infty,$$

that is  $(H^{n_k} \cdot M)_\infty \rightarrow (H \cdot M)_\infty$  in  $L^2$ . Hence as  $N$  is an  $L^2$  bounded martingale, the right hand side of the above expression tends to zero as  $k \rightarrow \infty$ . Similarly as  $(H^{n_k} \cdot N)_\infty \rightarrow (H \cdot N)_\infty$  in  $L^2$ , we see also that

$$\mathbb{E}((H^{n_k} \cdot N)_\infty M_\infty) \rightarrow 0, \text{ as } k \rightarrow \infty.$$

Hence we can pass to the limit to obtain the result for  $H$ .

To prove the second part of the theorem, we shall first show that

$$\langle (H \cdot N), (K \cdot M) \rangle + \langle (K \cdot N), (H \cdot M) \rangle = 2HK \langle M, N \rangle.$$

By polarisation

$$\langle M, N \rangle = \frac{\langle M + N \rangle - \langle M - N \rangle}{4},$$

also

$$HK = \frac{(H + K)^2 - (H - K)^2}{4}.$$

Hence

$$2(HK \cdot \langle M, N \rangle) = \frac{1}{8} \left( [(H + K)^2 - (H - K)^2] \cdot \{ \langle M + N \rangle - \langle M - N \rangle \} \right).$$

Now we use the result that  $\langle (H \cdot M) \rangle = (H^2 \cdot \langle M \rangle)$  which has been proved previously in theorem (7.9(iii)), to see that

$$\begin{aligned} 2(HK \cdot \langle M, N \rangle) &= \frac{1}{8} \left( \langle (H + K) \cdot (M + N) \rangle - \langle (H + K) \cdot (M - N) \rangle \right. \\ &\quad \left. - \langle (H - K) \cdot (M + N) \rangle + \langle (H - K) \cdot (M - N) \rangle \right). \end{aligned}$$

Considering the first two terms

$$\begin{aligned} \langle (H + K) \cdot (M + N) \rangle - \langle (H + K) \cdot (M - N) \rangle &= \\ &= \langle (H + K) \cdot M + (H + K) \cdot N \rangle - \langle (H + K) \cdot M - (H + K) \cdot N \rangle \\ &= 4 \langle (H + K) \cdot M, (H + K) \cdot N \rangle \quad \text{by polarisation} \\ &= 4 (\langle H \cdot M, H \cdot N \rangle + \langle H \cdot M, K \cdot N \rangle + \langle K \cdot M, H \cdot N \rangle + \langle K \cdot M, K \cdot N \rangle). \end{aligned}$$

Similarly for the second two terms

$$\begin{aligned} \langle (H - K) \cdot (M + N) \rangle - \langle (H - K) \cdot (M - N) \rangle &= \\ &= \langle (H - K) \cdot M + (H - K) \cdot N \rangle - \langle (H - K) \cdot M - (H - K) \cdot N \rangle \\ &= 4 \langle (H - K) \cdot M, (H - K) \cdot N \rangle \quad \text{by polarisation} \\ &= 4 (\langle H \cdot M, H \cdot N \rangle - \langle H \cdot M, K \cdot N \rangle - \langle K \cdot M, H \cdot N \rangle + \langle K \cdot M, K \cdot N \rangle). \end{aligned}$$

Adding these two together yields

$$2(HK \cdot \langle M, N \rangle) = \langle (H \cdot N), (K \cdot M) \rangle + \langle (K \cdot N), (H \cdot M) \rangle$$

Putting  $K \equiv 1$  yields

$$2H \cdot \langle M, N \rangle = \langle H \cdot M, N \rangle + \langle M, H \cdot N \rangle.$$

So it suffices to prove that  $\langle (H \cdot M), N \rangle = \langle M, (H \cdot N) \rangle$ , which is equivalent to showing that

$$(H \cdot M)N - (H \cdot N)M$$

is a local martingale (from the definition of covariation process). By localisation it suffices to consider  $M$  and  $N$  bounded martingales, whence we must check that for all stopping times  $T$ ,

$$\mathbb{E}((H \cdot M)_T N_T) = \mathbb{E}((H \cdot N)_T M_T),$$

but by the first part of the theorem

$$\mathbb{E}((H \cdot M)_\infty N_\infty) = \mathbb{E}((H \cdot N)_\infty M_\infty),$$

which is sufficient to establish the result, since

$$\begin{aligned} (H \cdot M)_T N_T &= (H \cdot M)_\infty^T N_\infty^T \\ (H \cdot N)_T M_T &= (H \cdot N)_\infty^T M_\infty^T \end{aligned}$$

□

**Corollary 11.2.**

Let  $N, M$  be continuous local martingales and  $H$  and  $K$  locally bounded previsible processes, then

$$\langle (H \cdot N), (K \cdot M) \rangle = (HK \cdot \langle N, M \rangle).$$

*Proof*

Note that the covariation is symmetric, hence

$$\begin{aligned} \langle (H \cdot N), (K \cdot M) \rangle &= (H \cdot \langle X, (K \cdot M) \rangle) \\ &= (H \cdot \langle (K \cdot M), X \rangle) \\ &= (HK \cdot \langle M, N \rangle). \end{aligned}$$

□

We can prove a stochastic calculus analogue of the usual integration by parts formula. However note that there is an extra term on the right hand side, the *covariation* of the processes  $X$  and  $Y$ . This is the first major difference we have seen between the Stochastic Integral and the usual Lebesgue Integral.

Before we can prove the general theorem, we need a lemma.

**Lemma (Parts for Finite Variation Process and a Martingale) 11.3.**

Let  $M$  be a bounded continuous martingale starting from zero, and  $V$  a bounded variation process starting from zero. Then

$$M_t V_t = \int_0^t M_s dV_s + \int_0^t V_s dM_s.$$

*Proof*

For  $n$  fixed, we can write

$$\begin{aligned} M_t V_t &= \sum_{k \geq 1} M_{k2^{-n} \wedge t} (V_{k2^{-n} \wedge t} - V_{(k-1)2^{-n} \wedge t}) + \sum_{k \geq 1} V_{(k-1)2^{-n} \wedge t} (M_{k2^{-n} \wedge t} - M_{(k-1)2^{-n} \wedge t}) \\ &= \sum_{k \geq 1} M_{k2^{-n} \wedge t} (V_{k2^{-n} \wedge t} - V_{(k-1)2^{-n} \wedge t}) + \int_0^t H_s^n dM_s, \end{aligned}$$

where  $H^n$  is the previsible simple process

$$H_s^n = \sum_{k \geq 1} V_{k2^{-n} \wedge t} 1_{((k-1)2^{-n} \wedge t, k2^{-n} \wedge t]}(s).$$

These  $H^n$  are bounded and converge to  $V$  by the continuity of  $V$ , so as  $n \rightarrow \infty$  the second term tends to

$$\int_0^t V_s dM_s,$$

and by the Dominated Convergence Theorem for Lebesgue-Stieltjes integrals, the second term converges to

$$\int_0^t M_s dV_s,$$

as  $n \rightarrow \infty$ . □

**Theorem (Integration by Parts) 11.4.**

For  $X$  and  $Y$  continuous semimartingales, then the following holds

$$X_t Y_t - X_0 Y_0 = \int_0^t X_s dY_s + \int_0^t Y_s dX_s + \langle X, Y \rangle_t.$$

*Proof*

It is trivial to see that it suffices to prove the result for processes starting from zero. Hence let  $X_t = M_t + A_t$  and  $Y_t = N_t + B_t$  in Doob-Meyer decomposition, so  $N_t$  and  $M_t$  are continuous local martingales and  $A_t$  and  $B_t$  are finite variation processes, all starting from zero. By localisation we can consider the local martingales  $M$  and  $N$  to be bounded martingales and the FV processes  $A$  and  $B$  to have bounded variation. Hence by the usual (finite variation) theory

$$A_t B_t = \int_0^t A_s dB_s + \int_0^t B_s dA_s.$$

It only remains to prove for bounded martingales  $N$  and  $M$  starting from zero that

$$M_t N_t = \int_0^t M_s dN_s + \int_0^t N_s dM_s + \langle M, N \rangle_t.$$

This follows by application of polarisation to corollary (7.3) to the quadratic variation existence theorem. Hence

$$\begin{aligned} (M_t + A_t)(N_t + B_t) &= M_t N_t + M_t B_t + N_t A_t + A_t B_t \\ &= \int_0^t M_s dN_s + \int_0^t N_s dM_s + \langle M, N \rangle_t \\ &\quad + \int_0^t M_s dB_s + \int_0^t B_s dM_s + \int_0^t N_s dA_s + \int_0^t A_s dN_s \\ &\quad + \int_0^t A_s dB_s + \int_0^t B_s dA_s \\ &= \int_0^t (M_s + A_s) d(N_s + B_s) + \int_0^t (N_s + B_s) d(M_s + A_s) + \langle M, N \rangle_t. \end{aligned}$$

□

Reflect for a moment that this theorem is implying another useful closure property of continuous semimartingales. It implies that the product of two continuous semimartingales  $X_t Y_t$  is a continuous semimartingale, since it can be written as a stochastic integrals with respect to continuous semimartingales and so it itself a continuous semimartingale.

**Theorem (Itô's Formula) 11.5.**

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a twice continuously differentiable function, and also let  $X = (X^1, X^2, \dots, X^n)$  be a continuous semimartingale in  $\mathbb{R}^n$ . Then

$$f(X_t) - f(X_0) = \sum_{i=1}^n \int_0^t \frac{\partial f}{\partial x^i}(X_s) dX_s^i + \frac{1}{2} \sum_{i,j=1}^n \int_0^t \frac{\partial^2 f}{\partial x^i \partial x^j}(X_s) d\langle X^i, X^j \rangle_s.$$

*Proof*

To prove Itô's formula; first consider the  $n = 1$  case to simplify the notation. Then let  $\mathcal{A}$  be the collection of  $C^2$  (twice differentiable) functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  for which it holds. Clearly  $\mathcal{A}$  is a vector space; in fact we shall show that it is also an algebra. To do this we must check that if  $f$  and  $g$  are in  $\mathcal{A}$ , then their product  $fg$  is also in  $\mathcal{A}$ . Let  $F_t = f(X_t)$  and  $G_t = g(X_t)$  be the associated semimartingales. From the integration by parts formula

$$F_t G_t - F_0 G_0 = \int_0^t F_s dG_s + \int_0^t G_s dF_s + \langle F_s, G_s \rangle_t.$$

However since by assumption  $f$  and  $g$  are in  $\mathcal{A}$ , Itô's formula may be applied to them individually, so

$$\int_0^t F_s dG_s = \int_0^t f(X_s) \frac{\partial f}{\partial x}(X_s) dX_s.$$

Also by the Kunita-Watanabe formula extended to continuous local martingales we have

$$\langle F, G \rangle_t = \int_0^t f'(X_s)g'(X_s)d\langle X, X \rangle_s.$$

Thus from the integration by parts,

$$\begin{aligned} F_t G_t - F_0 G_0 &= \int_0^t F_s dG_s + \int_0^t G_s dF_s + \int_0^t f'(X_s)g'(X_s)d\langle X, X \rangle_s, \\ &= \int_0^t (F_s g'(X_s) + f'(X_s)G_s) dX_s \\ &\quad + \frac{1}{2} \int_0^t (F_s g''(X_s) + 2f'(X_s)g'(X_s) + f''(X_s)G_s) d\langle M \rangle_s. \end{aligned}$$

So this is just what Itô's formula states for  $fg$  and so Itô's formula also applies to  $fg$ ; hence  $fg \in \mathcal{A}$ .

Since trivially  $f(x) = x$  is in  $\mathcal{A}$ , then as  $\mathcal{A}$  is an algebra, and a vector space this implies that  $\mathcal{A}$  contains all polynomials. So to complete the proof, we must approximate  $f$  by polynomials (which we can do by standard functional analysis), and check that in the limit we obtain Itô's formula.

Introduce a sequence  $U_n := \inf\{t : |X_t| + \langle X \rangle_t > n\}$ . Hence  $\{U_n\}$  is a sequence of stopping times tending to infinity. Now we shall prove Itô's formula for twice continuously differentiable  $f$  restricted to the interval  $[0, U_n]$ , so we can consider  $X$  as a bounded martingale. Consider a polynomial sequence  $f_k$  approximating  $f$ , in the sense that for  $r = 0, 1, 2$ ,  $f_k^{(r)} \rightarrow f^{(r)}$  uniformly on a compact interval. We have proved that Itô's formula holds for all polynomial, so it holds for  $f_k$  and hence

$$f_k(X_{t \wedge U_n}) - f_k(X_0) = \int_0^{t \wedge U_n} f'_k(X_s) dX_s + \frac{1}{2} \int_0^{t \wedge U_n} f''_k(X_s) d\langle X \rangle_s.$$

Let the continuous semimartingale  $X$  have Doob-Meyer decomposition

$$X_t = X_0 + M_t + A_t,$$

where  $M$  is a continuous local martingale and  $A$  is a finite variation process. We can rewrite the above as

$$f_k(X_{t \wedge U_n}) - f_k(X_0) = \int_0^{t \wedge U_n} f'_k(X_s) dM_s + \int_0^{t \wedge U_n} f'_k(X_s) dA_s + \frac{1}{2} \int_0^{t \wedge U_n} f''_k(X_s) d\langle M \rangle_s.$$

since  $\langle X \rangle = \langle M \rangle$ . On  $(0, U_n]$  the process  $|X|$  is uniformly bounded by  $n$ , so for  $r = 0, 1, 2$  from the convergence (which is uniform on the compact interval  $[0, U_n]$ ) we obtain

$$\sup_{|x| \leq n} |f_k^{(r)} - f^{(r)}| \rightarrow 0 \text{ as } k \rightarrow \infty$$

And from the Itô isometry we get the required convergence.

□

### 11.1. Applications of Itô's Formula

Let  $B_t$  be a standard Brownian motion; the aim of this example is to establish that:

$$\int_0^t B_s dB_s = \frac{1}{2} B_t^2 - \frac{1}{2} t.$$

This example gives a nice simple demonstration that all our hard work has achieved something. The result isn't the same as that which would be given by the 'logical' extension of the usual integration rules.

To prove this we apply Itô's formula to the function  $f(x) = x^2$ . We obtain

$$f(B_t) - f(B_0) = \int_0^t \frac{\partial f}{\partial x}(B_s) dB_s + \frac{1}{2} \int_0^t \frac{\partial^2 f}{\partial x^2}(B_s) d\langle B, B \rangle_s,$$

noting that  $B_0 = 0$  for a standard Brownian Motion we see that

$$B_t^2 = 2 \int_0^t B_s dB_s + \frac{1}{2} 2ds,$$

whence we derive that

$$\int_0^t B_s dB_s = \frac{B_t^2}{2} - \frac{t}{2}.$$

For those who have read the foregoing material carefully, there are grounds to complain that there are simpler ways to establish this result, notably by consideration of the definition of the quadratic variation process. However the point of this example was to show how Itô's formula can help in the actual evaluation of stochastic integrals; not to establish a totally new result.

### 11.2. Exponential Martingales

Exponential martingales play an important part in the theory. Suppose  $X$  is a continuous semimartingale starting from zero. Define:

$$Z_t = \exp\left(X_t - \frac{1}{2} \langle X \rangle_t\right).$$

This  $Z_t$  is called the **exponential semimartingale** associated with  $X_t$ , and it is the solution of the stochastic differential equation

$$dZ_t = Z_t dX_t,$$

that is

$$Z_t = 1 + \int_0^t Z_s dX_s,$$

so clearly if  $X$  is a continuous local martingale, i.e.  $X \in \mathcal{M}_{loc}^c$  then this implies, by the stability property of stochastic integration, that  $Z \in \mathcal{M}_{loc}^c$  also. In which case  $Z$  is the **exponential martingale**.

*Proof*

For existence, apply Itô's formula to  $f(x) = \exp(x)$  to obtain

$$d(\exp(Y_t)) = \exp(Y_t)dY_t + \frac{1}{2}\exp(Y_t)d\langle Y, Y \rangle_t.$$

Hence

$$\begin{aligned} d\left(\exp\left(X_t - \frac{1}{2}\langle X \rangle_t\right)\right) &= \exp\left(X_t - \frac{1}{2}\langle X \rangle_t\right)d\left(X_t - \frac{1}{2}\langle X \rangle_t\right) \\ &\quad + \frac{1}{2}\exp\left(X_t - \frac{1}{2}\langle X \rangle_t\right)d\left\langle X_t - \frac{1}{2}\langle X \rangle_t, X_t - \frac{1}{2}\langle X \rangle_t\right\rangle \\ &= \exp\left(X_t - \frac{1}{2}\langle X \rangle_t\right)dX_t - \frac{1}{2}\exp\left(X_t - \frac{1}{2}\langle X \rangle_t\right)d\langle X \rangle_t \\ &\quad + \frac{1}{2}\exp\left(X_t - \frac{1}{2}\langle X \rangle_t\right)d\langle X \rangle_t \\ &= Z_t dX_t \end{aligned}$$

Hence  $Z_t$  certainly solves the equation. Now to check uniqueness, define

$$Y_t = \exp\left(-X_t + \frac{1}{2}\langle X \rangle_t\right),$$

we wish to show that for every solution of the Stochastic Differential Equation  $Z_t Y_t$  is a constant. By a similar application of Itô's formula

$$dY_t = -Y_t dX_t + Y_t d\langle X \rangle_t,$$

whence by integration by parts (alternatively consider Itô applied to  $f(x, y) = xy$ ),

$$\begin{aligned} d(Z_t Y_t) &= Z_t dY_t + Y_t dZ_t + \langle Z, Y \rangle_t, \\ &= Z_t(-Y_t dX_t + Y_t d\langle X \rangle_t) + Y_t Z_t dX_t + (-Y_t Z_t)d\langle X \rangle_t, \\ &= 0. \end{aligned}$$

So  $Z_t Y_t$  is a constant, hence the unique solution of the stochastic differential equation  $dZ_t = Z_t dX_t$ , with  $Z_0 = 1$ , is

$$Z_t = \exp\left(X_t - \frac{1}{2}\langle X \rangle_t\right).$$

□

### Example

Let  $X_t = \theta B_t$ , for an arbitrary scalar  $\theta$ . Clearly  $X_t$  is a continuous local martingale, so the associated exponential martingale is

$$M_t = \exp\left(\theta B_t - \frac{1}{2}\theta^2 t\right).$$

## 12. Lévy Characterisation of Brownian Motion

---

A very useful result can be proved using the Itô calculus about the characterisation of Brownian Motion.

**Theorem 12.1.**

Let  $\{B^i\}_{t \geq 0}$  be continuous local martingales starting from zero for  $i = 1, \dots, n$ . Then  $B_t = (B_t^1, \dots, B_t^n)$  is a Brownian motion with respect to  $(\Omega, \mathcal{F}, \mathbb{P})$  adapted to the filtration  $\mathcal{F}_t$ , if and only iff

$$\langle B^i, B^j \rangle_t = \delta_{ij}t \quad \forall i, j \in \{1, \dots, n\}.$$

*Proof*

In these circumstances it follows that the statement  $B_t$  is a Brownian Motion is by definition equivalent to stating that  $B_t - B_s$  is independent of  $\mathcal{F}_s$  and is distributed normally with mean zero and covariance matrix  $(t - s)I$ .

Clearly if  $B_t$  is a Brownian motion then the covariation result follows trivially from the definitions. Now to establish the converse, we assume  $\langle B^i, B^j \rangle_t = \delta_{ij}t$  for  $i, j \in \{1, \dots, n\}$ , and shall prove  $B_t$  is a Brownian Motion.

Observe that for fixed  $\theta \in \mathbb{R}^n$  we can define  $M_t^\theta$  by

$$M_t^\theta := f(B_t, t) = \exp\left(i(\theta, x) + \frac{1}{2}|\theta|^2 t\right).$$

By application of Itô's formula to  $f$  we obtain (in differential form using the Einstein summation convention)

$$\begin{aligned} d(f(B_t, t)) &= \frac{\partial f}{\partial x^j}(B_t, t)dB_t^j + \frac{\partial f}{\partial t}(B_t, t)dt + \frac{1}{2} \frac{\partial^2 f}{\partial x^j \partial x^k}(B_t, t)d\langle B^j, B^k \rangle_t, \\ &= i\theta_j f(B_t, t)dB_t^j + \frac{1}{2}|\theta|^2 f(B_t, t)dt - \frac{1}{2}\theta_j \theta_k \delta_{jk} f(B_t, t)dt \\ &= i\theta_j f(B_t, t)dB_t^j. \end{aligned}$$

Hence

$$M_t^\theta = 1 + \int_0^t d(f(B_t, t)),$$

and is a sum of stochastic integrals with respect to continuous local martingales and is hence itself a continuous local martingale. But note that for each  $t$ ,

$$|M_t^\theta| = \left(e^{\frac{1}{2}|\theta|^2 t}\right) < \infty$$

Hence for any fixed time  $t_0$ ,  $(M^{t_0})_t$  satisfies

$$|(M^{t_0})_t| \leq |(M^{t_0})_\infty| < \infty,$$

[41]

and so is a bounded local martingale; hence  $(M^{t_0})_t$  is a martingale. Hence  $M^{t_0}$  is a genuine martingale. Thus for  $0 \leq s < t$  we have

$$\mathbb{E}(\exp(i(\theta, B_t - B_s)) | \mathcal{F}_s) = \exp\left(-\frac{1}{2}(t-s)|\theta|^2\right) \quad \text{a.s.}$$

However this is just the characteristic function of a normal random variable following  $N(0, t-s)$ ; so by the Lévy character theorem  $B_t - B_s$  is a  $N(0, t-s)$  random variable.  $\square$

## 13. Time Change of Brownian Motion

---

This result is one of frequent application, essentially it tells us that any continuous local martingale starting from zero, can be written as a time change of Brownian motion. So modulo a time change a Brownian motion is the most general kind of continuous local martingale.

**Proposition 13.1.**

Let  $M$  be a continuous local martingale starting from zero, such that  $\langle M \rangle_t \rightarrow \infty$  as  $t \rightarrow \infty$ . Then define

$$\tau_s := \inf\{t > 0 : \langle M \rangle_t > s\}.$$

Then define

$$\tilde{A}_s := M_{\tau_s}.$$

- (i) This  $\tau_s$  is an  $\mathcal{F}$  stopping time.
- (ii)  $\langle M \rangle_{\tau_s} = s$ .
- (iii) The local martingale  $M$  can be written as a time change of Brownian Motion as  $M_t = B_{\langle M \rangle_t}$ . Moreover the process  $\tilde{A}_s$  is an  $\tilde{\mathcal{F}}_s$  adapted Brownian Motion, where  $\tilde{\mathcal{F}}_s$  is the time-changed  $\sigma$  algebra i.e.  $\tilde{\mathcal{F}}_s = \mathcal{F}_{\tau_s}$ .

*Proof*

We may assume that the map  $t \mapsto \langle M \rangle_t$  is strictly increasing. Note that the map  $s \mapsto \tau_s$  is the inverse to  $t \mapsto \langle M \rangle_t$ . Hence the results (i),(ii) and (iii).

Define

$$\begin{aligned} T_n &:= \inf\{t : |M|_t > n\}, \\ [U_n &:= \langle M \rangle_{T_n}. \end{aligned}$$

Note that from these definitions

$$\begin{aligned} \tau_{t \wedge U_n} &= \inf\{s > 0 : \langle M \rangle_s > t \wedge U_n\} \\ &= \inf\{s > 0 : \langle M \rangle_s > t \wedge \langle M \rangle_{T_n}\} \\ &= T_n \wedge \tau_t \end{aligned}$$

So

$$\tilde{A}_s^{U_n} = \tilde{A}_{s \wedge U_n} = M_{\tau_t}^{T_n}.$$

Now note that  $U_n$  is an  $\tilde{\mathcal{F}}_t$  stopping time, since consider

$$\Lambda \equiv \{U_n \leq t\} \equiv \{\langle M \rangle_{T_n} \leq t\} \equiv \{T_n \leq \tau_t\},$$

the latter event is clearly  $\mathcal{F}_{\tau_t}$  measurable i.e. it is  $\tilde{\mathcal{F}}_t$  measurable, so  $U_n$  is a  $\tilde{\mathcal{F}}_t$  stopping time. We may now apply the optional stopping theorem to the UI martingale  $M^{T_n}$ , which yields

$$\begin{aligned} \mathbb{E}(\tilde{A}_t^{U_n} | \mathcal{F}_s) &= \mathbb{E}(\tilde{A}_{t \wedge U_n} | \tilde{\mathcal{F}}_s) = \mathbb{E}(M_{\tau_t}^{T_n} | \tilde{\mathcal{F}}_s) \\ &= \mathbb{E}(M_{\tau_t}^{T_n} | \mathcal{F}_{\tau_s}) = M_{\tau_s}^{T_n} = \tilde{A}_s^{U_n}. \end{aligned}$$

So  $\tilde{A}_t$  is a  $\tilde{\mathcal{F}}_t$  local martingale. But we also know that  $(M^2 - \langle M \rangle)^{T_n}$  is a UI martingale, since  $M^{T_n}$  is a UI martingale. By the optional stopping theorem, for  $0 < r < s$  we have

$$\begin{aligned} \mathbb{E} \left( \tilde{A}_{s \wedge U_n}^2 - (s \wedge U_n) | \tilde{\mathcal{F}}_r \right) &= \mathbb{E} \left( \left( (M_{\tau_s}^{T_n})^2 - \langle M \rangle_{\tau_s \wedge T_n} \right) | \mathcal{F}_{\tau_r} \right) \\ &= \mathbb{E} \left( (M_{\tau_s}^2 - \langle M \rangle_{\tau_s})^{T_n} | \mathcal{F}_{\tau_r} \right) = (M_{\tau_r}^2 - \langle M \rangle_{\tau_r})^{T_n} \\ &= \tilde{A}_{r \wedge U_n}^2 - (r \wedge U_n). \end{aligned}$$

Hence  $\tilde{A}^2 - t$  is a  $\tilde{\mathcal{F}}_t$  local martingale. Before we can apply Lévy's characterisation theorem we must check that  $\tilde{A}$  is continuous; that is we must check that for almost every  $\omega$  that  $M$  is constant on each interval of constancy of  $\langle M \rangle$ . By localisation it suffices to consider  $M$  a square integrable martingale, now let  $q$  be a positive rational, and define

$$S_q := \inf \{ t > q : \langle M \rangle_t > \langle M \rangle_q \},$$

then it is enough to show that  $M$  is constant on  $[q, S_q)$ . But  $M^2 - \langle M \rangle$  is a martingale, hence

$$\begin{aligned} \mathbb{E} \left[ \left( M_{S_q}^2 - \langle M \rangle_{S_q} \right)^2 | \mathcal{F}_q \right] &= M_q^2 - \langle M \rangle_q \\ &= M_q^2 - \langle M \rangle_{S_q}, \text{ as } \langle M \rangle_q = \langle M \rangle_{S_q}. \end{aligned}$$

Hence

$$\mathbb{E} \left[ (M_{S_q} - M_q)^2 | \mathcal{F}_q \right] = 0,$$

which establishes that  $\tilde{A}$  is continuous.

Thus  $\tilde{A}$  is a continuous  $\tilde{\mathcal{F}}_t$  adapted martingale with  $\langle \tilde{A} \rangle_s = s$  and so by the Lévy characterisation theorem  $\tilde{A}_s$  is a Brownian Motion.  $\square$

### 13.1. Gaussian Martingales

The time change of Brownian Motion can be used to prove the following useful theorem.

#### Theorem 13.2.

*If  $M$  is a continuous local martingale starting from zero, and  $\langle M \rangle_t$  is deterministic, that is if we can find a deterministic function  $f$  taking values in the non-negative real numbers such that  $\langle M \rangle_t = f(t)$  a.e., then  $M$  is a Gaussian Martingale (i.e.  $M_t$  has a Gaussian distribution for almost all  $t$ ).*

*Proof*

Note that by the time change of Brownian Motion theorem, we can write  $M_t$  as a time change of Brownian Motion through

$$M_t = B_{\langle M \rangle_t},$$

where  $B$  is a standard Brownian Motion. By hypothesis  $\langle M \rangle_t = f(t)$ , a deterministic function for almost all  $t$ , hence for almost all  $t$ ,

$$M_t = B_{f(t)},$$

but the right hand side is a gaussian random variable following  $N(0, f(t))$ . Hence  $M$  is a Gaussian Martingale, and at time  $t$  it has distribution given by  $N(0, \langle M \rangle_t)$ .  $\square$

As a corollary consider the stochastic integral of a purely deterministic function with respect to a brownian motion.

**Corollary 13.3.**

Let  $g(t)$  be a deterministic function of  $t$ , then  $M$  defined by

$$M_t := \int_0^t f(s)dB_s,$$

satisfies

$$M_t \sim N\left(0, \int_0^t |f(s)|^2 ds\right).$$

*Proof*

From the definition of  $M$  via a stochastic integral with respect to a continuous martingale, it is clear that  $M$  is a continuous local martingale, and by the Kunita-Watanabe result, the quadratic variation of  $M$  is given by

$$\langle M \rangle_t = \int_0^t |f(s)|^2 ds,$$

hence the result follows.  $\square$

This result can also be established directly in a fashion which is very similar to the proof of the Lévy characterisation theorem. Consider  $Z$  defined via

$$Z_t = \exp\left(i\theta M_t + \frac{1}{2}\theta^2 \langle M \rangle_t\right),$$

as in the Lévy characterisation proof, we see that this is a continuous local martingale, and by boundedness furthermore is a martingale, and hence

$$\mathbb{E}(Z_0) = \mathbb{E}(Z_t),$$

whence

$$\mathbb{E}(\exp(i\theta M_t)) = \mathbb{E}\left(\exp\left(-\frac{1}{2}\theta^2 \int_0^t |f(s)|^2 ds\right)\right)$$

which is the characteristic function of the appropriate normal distribution.

## 14. Girsanov's Theorem

---

Girsanov's theorem is an element of stochastic calculus which does not have an analogue in standard calculus.

### 14.1. Change of measure

When we wish to compare two measures  $\mathbb{P}$  and  $\mathbb{Q}$ , we don't want either of them simply to throw information away; since when they are positive they can be related by the *Radon-Nikodym* derivative; this motivates the following definition of *equivalence* of two measures.

**Definition 14.1.**

Two measures  $\mathbb{P}$  and  $\mathbb{Q}$  are said to be *equivalent* if they operate on the same sample space, and if  $A$  is any event in the sample space then

$$\mathbb{P}(A) > 0 \Leftrightarrow \mathbb{Q}(A) > 0.$$

In other words  $\mathbb{P}$  is absolutely continuous with respect to  $\mathbb{Q}$  and  $\mathbb{Q}$  is absolutely continuous with respect to  $\mathbb{P}$ .

**Theorem 14.2.**

If  $\mathbb{P}$  and  $\mathbb{Q}$  are equivalent measures, and  $X_t$  is an  $\mathcal{F}_t$ -adapted process then the following results hold

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}(X_t) &= \mathbb{E}_{\mathbb{P}}\left(\frac{d\mathbb{Q}}{d\mathbb{P}}X_t\right), \\ \mathbb{E}_{\mathbb{Q}}(X_t|\mathcal{F}_s) &= L_s^{-1}\mathbb{E}_{\mathbb{P}}(L_tX_t|\mathcal{F}_s), \end{aligned}$$

where

$$L_s = \mathbb{E}_{\mathbb{P}}\left(\frac{d\mathbb{Q}}{d\mathbb{P}}\middle|\mathcal{F}_s\right).$$

Here  $L_t$  is the Radon-Nikodym derivative of  $\mathbb{Q}$  with respect to  $\mathbb{P}$ . The first result basically shows that this is a martingale, and the second is a continuous time version of Bayes theorem.

*Proof*

The first part is basically the statement that the Radon-Nikodym derivative is a martingale. This follows because the measures  $\mathbb{P}$  and  $\mathbb{Q}$  are equivalent, but this will not be proved in detail here. Let  $Y$  be an  $\mathcal{F}_t$  measurable random variable, such that  $\mathbb{E}_{\mathbb{Q}}(|Y|) < \infty$ . We shall prove that

$$\mathbb{E}_{\mathbb{Q}}(Y|\mathcal{F}_s) = \frac{1}{L_s}\mathbb{E}_{\mathbb{P}}[YL_t|\mathcal{F}_s] \text{ a.s. } (\mathbb{P} \text{ and } \mathbb{Q}).$$

Then for any  $A \in \mathcal{F}_s$ , using the definition of conditional expectation we have that

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}}\left(1_A \frac{1}{L_s}\mathbb{E}_{\mathbb{P}}[YL_t|\mathcal{F}_s]\right) &= \mathbb{E}_{\mathbb{P}}(1_A\mathbb{E}_{\mathbb{P}}[YL_t|\mathcal{F}_s]) \\ &= \mathbb{E}_{\mathbb{P}}[1_AYL_t] = \mathbb{E}_{\mathbb{Q}}[1_AY]. \end{aligned}$$

Substituting  $Y = X_t$  gives the desired result. □

**Theorem (Girsanov).**

Let  $M$  be a continuous local martingale, and let  $Z$  be the associated exponential martingale

$$Z_t = \exp\left(M_t - \frac{1}{2}\langle M \rangle_t\right).$$

If  $Z$  is uniformly integrable, then a new measure  $\mathbb{Q}$ , equivalent to  $\mathbb{P}$  may be defined by

$$\frac{d\mathbb{Q}}{d\mathbb{P}} = Z_\infty.$$

Then if  $X$  is a continuous  $\mathbb{P}$  local martingale,  $X - \langle X, M \rangle$  is a  $\mathbb{Q}$  local martingale.

*Proof*

Since  $Z_\infty$  exists a.s. it defines a uniformly integrable martingale (the exponential martingale), a version of which is given by  $Z_t = \mathbb{E}(Z_\infty | \mathcal{F}_t)$ . Hence  $\mathbb{Q}$  constructed thus is a probability measure which is equivalent to  $\mathbb{P}$ . Now consider  $X$ , a  $\mathbb{P}$  local martingale. Define a sequence of stopping times which tend to infinity via

$$T_n := \inf\{t \geq 0 : |X_t| \geq n, \text{ or } |\langle X, M \rangle_t| \geq n\}.$$

Now consider the process  $Y$  defined via

$$Y := X^{T_n} - \langle X^{T_n}, M \rangle.$$

By Itô's formula for  $0 \leq t \leq T_n$ , remembering that  $dZ_t = Z_t dM_t$  as  $Z$  is the exponential martingale associated with  $M$ ,

$$\begin{aligned} d(Z_t Y_t) &= Z_t dY_t + Y_t dZ_t + \langle Z, Y \rangle \\ &= Z_t (dX_t - d\langle X, M \rangle) + Y_t Z_t dM_t + \langle Z, Y \rangle \\ &= Z_t (dX_t - d\langle X, M \rangle) + (X_t - \langle X, M \rangle_t) Z_t dM_t + Z_t d\langle X, M \rangle \\ &= (X_t - \langle X, M \rangle_t) Z_t dM_t + Z_t dX_t \end{aligned}$$

Where the result  $\langle Z, Y \rangle_t = Z_t \langle X, M \rangle_t$  follows from the Kunita-Watanabe theorem. Hence  $ZY$  is a  $\mathbb{P}$ -local martingale. But since  $Z$  is uniformly integrable, and  $Y$  is bounded (by construction of the stopping time  $T_n$ ), hence  $ZY$  is a genuine  $\mathbb{P}$ -martingale. Hence for  $s < t$  and  $A \in \mathcal{F}_s$ , we have

$$\mathbb{E}_{\mathbb{Q}}[(Y_t - Y_s)1_A] = \mathbb{E}[Z_\infty(Y_t - Y_s)1_A] = \mathbb{E}[(Z_t Y_t - Z_s Y_s)1_A] = 0,$$

hence  $Y$  is a  $\mathbb{Q}$  martingale. Thus  $X - \langle X, M \rangle$  is a  $\mathbb{Q}$  local martingale, since  $T_n$  is a reducing sequence such that  $(X - \langle X, M \rangle)^{T_n}$  is a  $\mathbb{Q}$ -martingale, and  $T_n \uparrow \infty$  as  $n \rightarrow \infty$ .  $\square$

**Corollary 14.3.**

Let  $W_t$  be a  $\mathbb{P}$  Brownian motion, then  $\tilde{W}_t := W_t - \langle W, M \rangle_t$  is a  $\mathbb{Q}$  Brownian motion.

*Proof*

Use Lévy's characterisation of Brownian motion to see that since  $\tilde{W}_t$  is continuous and  $\langle \tilde{W}, \tilde{W} \rangle_t = \langle W, W \rangle_t = t$ , since  $W_t$  is a  $\mathbb{P}$  Brownian motion, then  $\tilde{W}$  is a  $\mathbb{Q}$  Brownian motion.  $\square$

## 15. Relations to Second Order PDEs

---

The aim of this section is to show a rather surprising connection between stochastic differential equations and the solution of second order partial differential equations. Surprising though the results may seem they often provide a viable route to calculating the solutions of explicit PDEs (an example of this is solving the Black-Scholes Equation in Option Pricing, which is much easier to solve via stochastic methods, than as a second order PDE). At first this may well seem to be surprising since one problem is entirely deterministic and the other is inherently stochastic!

### 15.1. Infinitesimal Generator

Consider the following  $d$ -dimensional SDE,

$$\begin{aligned} d\mathbf{X}_t &= \mathbf{b}(\mathbf{X}_t)dt + \sigma(\mathbf{X}_t)d\mathbf{B}_t, \\ \mathbf{X}_0 &= x_0 \end{aligned}$$

where  $\sigma$  is a  $d \times d$  matrix with elements  $\sigma = \{\sigma_{ij}\}$ . This SDE has infinitesimal generator  $A$ , where

$$A = \sum_{j=1}^d b^j(\mathbf{X}_t) \frac{\partial}{\partial x^j} + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d \sum_{k=1}^d \sigma_{ik}(\mathbf{X}_t) \sigma_{kj}(\mathbf{X}_t) \frac{\partial^2}{\partial x^i \partial x^j}.$$

It is conventional to set

$$a_{ij} = \sum_{k=1}^d \sigma_{ik} \sigma_{kj},$$

whence  $A$  takes the simpler form

$$A = \sum_{j=1}^d b^j(\mathbf{X}_t) \frac{\partial}{\partial x^j} + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d a_{ij}(\mathbf{X}_t) \frac{\partial^2}{\partial x^i \partial x^j}.$$

Why is the definition useful? Consider application of Itô's formula to  $f(\mathbf{X}_t)$ , which yields

$$f(\mathbf{X}_t) - f(\mathbf{X}_0) = \int_0^t \sum_{j=1}^d \frac{\partial f}{\partial x^j}(\mathbf{X}_s) d\mathbf{X}_s + \frac{1}{2} \int_0^t \sum_{i=1}^d \sum_{j=1}^d \frac{\partial^2 f}{\partial x^i \partial x^j}(\mathbf{X}_s) d\langle X^i, X^j \rangle_s.$$

Substituting for  $d\mathbf{X}_t$  from the SDE we obtain,

$$\begin{aligned} f(\mathbf{X}_t) - f(\mathbf{X}_0) &= \int_0^t \left( \sum_{j=1}^d b^j(\mathbf{X}_s) \frac{\partial f}{\partial x^j}(\mathbf{X}_s) + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d \sum_{k=1}^d \sigma_{ik} \sigma_{kj} \frac{\partial^2 f}{\partial x^i \partial x^j}(\mathbf{X}_s) \right) dt \\ &\quad + \int_0^t \sum_{j=1}^d \sigma_{ij}(\mathbf{X}_s) \frac{\partial f}{\partial x^j}(\mathbf{X}_s) d\mathbf{B}_s \\ &= \int_0^t Af(\mathbf{X}_s) ds + \int_0^t \sum_{j=1}^d \sigma_{ij}(\mathbf{X}_s) \frac{\partial f}{\partial x^j}(\mathbf{X}_s) d\mathbf{B}_s \end{aligned}$$

[48]

**Definition 15.1.**

We say that  $X_t$  satisfies the martingale problem for  $A$ , if  $X_t$  is  $\mathcal{F}_t$  adapted and

$$M_t = f(\mathbf{X}_t) - f(\mathbf{X}_0) - \int_0^t Af(X_s)ds,$$

is a martingale for each  $f \in C_c^2(\mathbb{R}^d)$ .

It is simple to verify from the foregoing that any solution of the associated SDE solves the martingale problem for  $A$ . This can be generalised if we consider test functions  $\phi \in C^2(\mathbb{R}^+ \times \mathbb{R}^d, \mathbb{R})$ , and define

$$M_t^\phi := \phi(t, \mathbf{X}_t) - \phi(0, \mathbf{X}_0) - \int_0^t \left( \frac{\partial}{\partial s} + A \right) \phi(s, \mathbf{X}_s) ds.$$

then  $M_t^\phi$  is a local martingale, for  $\mathbf{X}_t$  a solution of the SDE associated with the infinitesimal generator  $A$ . The proof follows by an application of Itô's formula to  $M_t^\phi$ , similar to that of the above discussion.

**15.2. The Dirichlet Problem**

Let  $\Omega$  be a subspace of  $\mathbb{R}^d$  with a smooth boundary  $\partial\Omega$ . The *Dirichlet Problem* for  $f$  is defined as the solution of the system

$$\begin{aligned} Au + \phi &= 0 \text{ on } \Omega, \\ u &= f \text{ on } \partial\Omega. \end{aligned}$$

Where  $A$  is a second order partial differential operator of the form

$$A = \sum_{j=1}^d b^j(\mathbf{X}_t) \frac{\partial}{\partial x^j} + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d a_{ij}(\mathbf{X}_t) \frac{\partial^2}{\partial x^i \partial x^j},$$

which is associated as before to an SDE. This SDE will play an important role in what is to follow.

A simple example of a Dirichlet Problem is the solution of the Laplace equation in the disc, with Dirichlet boundary conditions on the boundary, i.e.

$$\begin{aligned} \nabla^2 u &= 0 \text{ on } D, \\ u &= f \text{ on } \partial D. \end{aligned}$$

**Theorem 15.2.**

For each  $f \in C_b^2(\partial\Omega)$  there exists a unique  $u \in C_b^2(\bar{\Omega})$  solving the Dirichlet problem for  $f$ . Moreover there exists a continuous function  $m : \bar{\Omega} \rightarrow (0, \infty)$  such that for all  $f \in C_b^2(\partial\Omega)$  this solution is given by

$$u(\mathbf{x}) = \int_{\partial\Omega} m(\mathbf{x}, \mathbf{y}) f(\mathbf{y}) \sigma(d\mathbf{y}).$$

Now remember the SDE which is associated with the infinitesimal generator  $A$ :

$$\begin{aligned} d\mathbf{X}_t &= \mathbf{b}(\mathbf{X}_t)dt + \sigma(\mathbf{X}_t)d\mathbf{B}_t, \\ \mathbf{X}_0 &= x_0 \end{aligned}$$

Often in what follows we shall want to consider the conditional expectation and probability measures, conditional on  $x_0 = x$ , these will be denoted  $\mathbb{E}_x$  and  $\mathbb{P}_x$  respectively.

**Theorem (Dirichlet Solution).**

Define a stopping time via

$$T := \inf\{t \geq 0 : X_t \notin \Omega\}.$$

Then  $u(\mathbf{x})$  given by

$$u(\mathbf{x}) := \mathbb{E}_x \left[ \int_0^T \phi(\mathbf{X}_s)ds + f(\mathbf{X}_T) \right],$$

solves the Dirichlet problem for  $f$ .

*Proof*

Define

$$M_t := u(\mathbf{X}_{T \wedge t}) + \int_0^{t \wedge T} \phi(\mathbf{X}_s)ds.$$

We shall now show that this  $M_t$  is a martingale. For  $t \geq T$ , it is clear that  $dM_t = 0$ . For  $t < T$  by Itô's formula

$$dM_t = du(\mathbf{X}_t) + \phi(\mathbf{X}_t)dt.$$

Also, by Itô's formula,

$$\begin{aligned} du(\mathbf{X}_t) &= \sum_{j=1}^d \frac{\partial u}{\partial x^j}(\mathbf{X}_t) dX_t^j + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d \frac{\partial^2 u}{\partial x^i \partial x^j}(\mathbf{X}_t) d\langle X^i, X^j \rangle_t \\ &= \sum_{j=1}^d \frac{\partial u}{\partial x^j}(\mathbf{X}_t) [\mathbf{b}(\mathbf{X}_t)dt + \sigma(\mathbf{X}_t)d\mathbf{B}_t] + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d \sum_{k=1}^d \sigma_{ik}\sigma_{kj} \frac{\partial^2 u}{\partial x^i \partial x^j}(\mathbf{X}_t)dt \\ &= Au(\mathbf{X}_t)dt + \sum_{j=1}^d \sigma(\mathbf{X}_t) \frac{\partial u}{\partial x^j}(\mathbf{X}_t) d\mathbf{B}_t. \end{aligned}$$

Putting these two applications of Itô's formula together yields

$$dM_t = (Au(\mathbf{X}_t) + \phi(\mathbf{X}_t)) dt + \sum_{j=1}^d \sigma(\mathbf{X}_t) \frac{\partial u}{\partial x^j}(\mathbf{X}_t) d\mathbf{B}_t.$$

but since  $u$  solves the Dirichlet problem, then

$$(Au + \phi)(\mathbf{X}_t) = 0,$$

hence

$$\begin{aligned} dM_t &= (Au(\mathbf{X}_t) + \phi(\mathbf{X}_t)) dt + \sum_{j=1}^d \sigma(\mathbf{X}_t) \frac{\partial u}{\partial x^j}(\mathbf{X}_t) d\mathbf{B}_t^j, \\ &= \sum_{j=1}^d \sigma(\mathbf{X}_t) \frac{\partial u}{\partial x^j}(\mathbf{X}_t) d\mathbf{B}_t^j. \end{aligned}$$

from which we conclude by the stability property of the stochastic integral that  $M_t$  is a local martingale. However  $M_t$  is uniformly bounded on  $[0, t]$ , and hence  $M_t$  is a martingale.

In particular, let  $\phi(x) \equiv 1$ , and  $f \equiv 0$ , by the optional stopping theorem, since  $T \wedge t$  is a bounded stopping time, this gives

$$u(\mathbf{x}) = \mathbb{E}_x(M_0) = \mathbb{E}_x(M_{T \wedge t}) = \mathbb{E}_x[u(\mathbf{X}_{T \wedge t}) + (T \wedge t)].$$

Letting  $t \rightarrow \infty$ , we have via monotone convergence that  $\mathbb{E}_x(T) < \infty$ , since we know that the solutions  $u$  is bounded from the PDE solution existence theorem; hence  $T < \infty$  a.s.. We cannot simply apply the optional stopping theorem directly, since  $T$  is not necessarily a bounded stopping time. But for arbitrary  $\phi$  and  $f$ , we have that

$$|M_t| \leq \|u\|_\infty + T\|\phi\|_\infty = \sup_{\mathbf{x} \in \bar{\Omega}} |u(\mathbf{x})| + T \sup_{\mathbf{x} \in \bar{\Omega}} |\phi(\mathbf{x})|,$$

whence as  $\mathbb{E}_x(T) < \infty$ , the martingale  $M$  is uniformly integrable, and by the martingale convergence theorem has a limit  $M_\infty$ . This limiting random variable is given by

$$M_\infty = f(\mathbf{X}_T) + \int_0^T \phi(\mathbf{X}_s) ds.$$

Hence from the identity  $\mathbb{E}_x M_0 = \mathbb{E}_x M_\infty$  we have that,

$$u(\mathbf{x}) = \mathbb{E}_x(M_0) = \mathbb{E}_x(M_\infty) = \mathbb{E}_x \left[ f(\mathbf{X}_T) + \int_0^T \phi(\mathbf{X}_s) ds \right].$$

□

### 15.3. The Cauchy Problem

The *Cauchy Problem* for  $f$ , a  $C_b^2$  function, is the solution of the system

$$\begin{aligned} \frac{\partial u}{\partial t} &= Au \text{ on } \Omega \\ u(0, \mathbf{x}) &= f(\mathbf{x}) \text{ on } \mathbf{x} \in \Omega \\ u(t, \mathbf{x}) &= f(\mathbf{x}) \quad \forall t \geq 0, \text{ on } \mathbf{x} \in \partial\Omega \end{aligned}$$

A typical problem of this sort is to solve the heat equation,

$$\frac{\partial u}{\partial t} = \frac{1}{2} \nabla^2 u,$$

where the function  $u$  represents the temperature in a region  $\Omega$ , and the boundary condition is to specify the temperature field over the region at time zero, i.e. a condition of the form

$$u(0, \mathbf{x}) = f(\mathbf{x}) \text{ for } x \in \Omega,$$

In addition the boundary has its temperature fixed at zero,

$$u(0, \mathbf{x}) = 0 \text{ for } x \in \partial\Omega.$$

If  $\Omega$  is just the real line, then the solution has the beautifully simple form

$$u(t, x) = \mathbb{E}_x (f(B_t)),$$

where  $B_t$  is a standard Brownian Motion.

**Theorem (Cauchy Existence) 15.3.**

For each  $f \in C_b^{1,2}(\mathbb{R}^d)$  there exists a unique  $u$  in  $C_b^{1,2}(\mathbb{R} \times \mathbb{R}^d)$  such that  $u$  solves the Cauchy Problem for  $f$ . Moreover there exists a continuous function (the heat kernel)

$$p : (0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d \rightarrow (0, \infty),$$

such that for all  $f \in C_b^2(\mathbb{R}^d)$ , the solution to the Cauchy Problem for  $f$  is given by

$$u(t, \mathbf{x}) = \int_{\mathbb{R}^d} p(t, \mathbf{x}, \mathbf{y}) f(\mathbf{y}) d\mathbf{y}.$$

**Theorem 15.4.**

Let  $u \in C_b^{1,2}(\mathbb{R} \times \mathbb{R}^d)$  be the solution of the Cauchy Problem for  $f$ . Then define

$$T := \inf\{t \geq 0 : \mathbf{X}_t \notin \Omega\},$$

a stopping time. Then

$$u(t, \mathbf{x}) = \mathbb{E}_x [f(\mathbf{X}_{T \wedge t})]$$

*Proof*

Fix  $s \in (0, \infty)$  and consider the time reversed process

$$M_t := u((s-t) \wedge T, \mathbf{X}_{t \wedge T}).$$

There are three cases now to consider; for  $0 \leq T \leq t \leq s$ ,  $M_t = u((s-t) \wedge T, \mathbf{X}_T)$ , where  $X_T \in \partial\Omega$ , so from the boundary condition,  $M_t = f(\mathbf{X}_T)$ , and hence it is clear that  $dM_t = 0$ . For  $0 \leq s \leq T \leq t$  and for  $0 \leq t \leq s \leq T$ , the argument is similar; in the latter case by Itô's formula we obtain

$$\begin{aligned} dM_t &= -\frac{\partial u}{\partial t}(s-t, \mathbf{X}_t)dt + \sum_{j=1}^d \frac{\partial u}{\partial x^j}(s-t, \mathbf{X}_t)dX_t^j \\ &\quad + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d \frac{\partial^2 u}{\partial x^i \partial x^j}(s-t, \mathbf{X}_t)d\langle X^i, X^j \rangle_t, \\ &= \left( -\frac{\partial u}{\partial t} + Au \right) (s-t, \mathbf{X}_t)dt + \sum_{j=1}^d \frac{\partial u}{\partial x^j}(s-t, \mathbf{X}_t) \sum_{k=1}^d \sigma_{jk}(\mathbf{X}_t) d\mathbf{B}_t^k. \end{aligned}$$

We obtain a similar result in the  $0 \leq t \leq T \leq s$ , case but with  $s$  replaced by  $T$ . Thus for  $u$  solving the Cauchy Problem for  $f$ , we have that

$$\left(-\frac{\partial u}{\partial t} + Au\right) = 0,$$

we see that  $M_t$  is a local martingale. Boundedness implies that  $M_t$  is a martingale, and hence by optional stopping

$$u(s, \mathbf{x}) = \mathbb{E}_x(M_0) = \mathbb{E}_x(M_s) = \mathbb{E}_x(f(\mathbf{X}_{s \wedge T})),$$

□

### 15.4. Feynman-Kač Representation

Feynman observed the following representation for the representation of the solution of a PDE via the expectation of a suitable function of a Brownian Motion ‘intuitively’ and the theory was later made rigorous by Kač.

We have already considered solving the Cauchy problem

$$\begin{aligned} \frac{\partial u}{\partial t} &= Au \text{ on } \Omega \\ u(0, \mathbf{x}) &= f(\mathbf{x}) \text{ on } \mathbf{x} \in \Omega \\ u(t, \mathbf{x}) &= f(\mathbf{x}) \forall t \geq 0, \text{ on } \mathbf{x} \in \partial\Omega \end{aligned}$$

where  $A$  is the generator of an SDE and hence of the form

$$A = \sum_{j=1}^d b^j(\mathbf{X}_t) \frac{\partial}{\partial x^j} + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d a_{ij}(\mathbf{X}_t) \frac{\partial^2}{\partial x^i \partial x^j}.$$

Now consider the more general form of the same Cauchy problem where we consider a Cauchy Problem with generator  $L$  of the form:

$$L \equiv A + v = \sum_{j=1}^d b^j(\mathbf{X}_t) \frac{\partial}{\partial x^j} + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d a_{ij}(\mathbf{X}_t) \frac{\partial^2}{\partial x^i \partial x^j} + v(\mathbf{X}_t).$$

For example

$$A = \frac{1}{2} \nabla^2 + v(\mathbf{X}_t),$$

so in this example we are solving the problem

$$\begin{aligned} \frac{\partial u}{\partial t} &= \frac{1}{2} \nabla^2 u(t, \mathbf{X}_t) + v(\mathbf{X}_t) u(\mathbf{X}_t) \text{ on } \mathbb{R}^d. \\ &\& u(0, \mathbf{x}) = f(\mathbf{x}) \text{ on } \partial\mathbb{R}^d. \end{aligned}$$

The Feynman-Kač Representation Theorem expresses the solution of a general second order PDE in terms of an expectation of a function of a Brownian Motion. To simplify the statement of the result, we shall work on  $\Omega = \mathbb{R}^d$ , since this removes the problem of considering the Brownian Motion hitting the boundary.

**Theorem (Feynman-Kač Representation).**

Let  $u \in C_b^{1,2}(\mathbb{R} \times \mathbb{R}^d)$  be a solution of the Cauchy Problem with a generator of the above form for  $f$ , and let  $\mathbf{B}_t$  be a Brownian Motion in  $\mathbb{R}^d$  starting at  $\mathbf{x}$ . Then

$$u(t, \mathbf{x}) = \mathbb{E}_x \left[ f(\mathbf{B}_t) \exp \left( \int_0^t v(\mathbf{B}_s) ds \right) \right].$$

*Proof*

Fix  $s \in (0, \infty)$  and apply Itô's formula to

$$M_t = u(s - t, \mathbf{X}_t) \exp \left( \int_0^t v(\mathbf{B}_r) dr \right).$$

For notational convenience, let

$$E_t = \exp \left( \int_0^t v(\mathbf{B}_r) dr \right).$$

For  $0 \leq t \leq s$ , we have

$$\begin{aligned} dM_t &= \sum_{j=1}^d \frac{\partial u}{\partial x^j}(s - t, \mathbf{X}_t) E_t dB_t^j + \left( \frac{\partial u}{\partial t} + Au + vu \right) (s - t, \mathbf{X}_t) E_t dt \\ &= \sum_{j=1}^d \frac{\partial u}{\partial x^j}(s - t, \mathbf{X}_t) E_t dB_t^j. \end{aligned}$$

Hence  $M_t$  is a local martingale; since it is bounded,  $M_t$  is a martingale and hence by optional stopping

$$u(s, \mathbf{x}) = \mathbb{E}_x(M_0) = \mathbb{E}_x(M_s) = \mathbb{E}_x(f(\mathbf{X}_s)E_s).$$

□

## 16. Stochastic Filtering

---

The aim of the stochastic filtering problem is to determine the law of the signal given the observations. This must be formalised in the language of stochastic calculus. The notation  $\|\cdot\|$  will be used in this section to denote the Euclidean norm in  $d$ -dimensional space.

### 16.1. Signal Process

Let  $\{X_t, \mathcal{F}_t, t \geq 0\}$  be the *signal process*. It is defined to be the solution of the stochastic differential equation

$$dX_t = f(t, X_t)dt + \sigma(t, X_t)dV_t,$$

where  $V$  is a  $d$  dimensional standard Brownian motion. The coefficients satisfy the conditions

$$\begin{aligned} \|f(t, x) - f(t, y)\| + \|\sigma(t, x) - \sigma(t, y)\| &\leq k\|x - y\| \\ \|f(t, x)\|^2 + \|\sigma(t, x)\|^2 &\leq k^2(1 + \|x\|^2), \end{aligned}$$

which ensure that the equation for  $X$  has a unique solution.

### 16.2. Observation Process

The observation process satisfies the stochastic differential equation

$$dY_t = h(t, X_t)dt + dW_t, \quad Y_0 = 0,$$

where  $W_t$  is an  $m$  dimensional standard Brownian motion. The function  $h$  satisfies a linear growth condition

$$\|h(t, x)\|^2 \leq k(1 + \|x\|^2).$$

A consequence of this condition is that for any  $T > 0$ ,

$$\mathbb{E}(\|X_t\|^2) \leq C(1 + \mathbb{E}[\|X_0\|])e^{ct},$$

for  $t \in [0, T]$  with suitable constants  $C$  and  $c$  which may be functions of  $T$ . As a result of this bound

$$\mathbb{E} \left[ \int_0^T |h(s, X_s)|^2 ds \right] < \infty.$$

Given this process a sequence of  $\sigma$ -algebras may be defined

$$\mathcal{Y}_t := \sigma(Y_s : 0 \leq s \leq t) \cup \mathcal{N},$$

where  $\mathcal{N}$  is the set of  $\mathbb{P}$  null subsets of  $\Omega$ , and

$$\mathcal{Y} := \bigcup_{t \geq 0} \mathcal{Y}_t.$$

### 16.3. The Filtering Problem

The above has set the scene, we have a real physical system whose state at time  $t$  is represented by the vector  $X_t$ . The system state is governed by an SDE with a noise term representing random perturbations to the system. This is observed to give the observation process  $Y_t$  which includes new independent noise (represented by  $W$ ). An example might be a GPS system on a rapidly moving object, where periodic position measurements may be obtained by the GPS system. The object obeys Newton's laws of motion (giving the signal SDE), and the observations are those of the GPS system (with inherent errors). The filtering problem is to find the conditional law of the signal process given the observations, i.e. to find

$$\pi_t(\phi) := \mathbb{E}(\phi(X_t)|\mathcal{Y}_t) \quad \forall t \in [0, T].$$

In the case mentioned above the filtering problem is to work out the position of the object given the sequence of GPS position measurements.

### 16.4. Change of Measure

To solve the filtering problem Girsanov's theorem will be used to make a change of measure to a new measure under which the observation process  $Y$  is a Brownian motion. Let

$$Z_t := \exp\left(-\int_0^t h^T(s, X_s)dW_s - \frac{1}{2}\int_0^t |h(s, X_s)|^2 ds\right).$$

#### Proposition 16.1.

$\{Z_t, \mathcal{F}_t, t \geq 0\}$  is a martingale.

*Proof*

The process  $Z_t$  satisfies

$$Z_t = 1 - \int_0^t Z_s h^T(s, X_s)dW_s,$$

that is  $Z_t$  solves the SDE

$$\begin{aligned} dZ_t &= -Z_t h^T(t, X_t)dW_t, \\ Z_0 &= 1. \end{aligned}$$

Given this expression,  $Z_t$  is a positive continuous local martingale and hence is a supermartingale. To prove that it is a true martingale we must prove in addition that it has constant mean.

Let  $T_n$  be a reducing sequence for the local martingale  $Z_t$ , i.e. an increasing sequence of stopping times tending to infinity as  $n \rightarrow \infty$  such that  $Z^{T_n}$  is a genuine martingale. By Fatou's lemma, and the local martingale property

$$\mathbb{E}Z_t = \mathbb{E} \lim_{n \rightarrow \infty} Z_t^{T_n} \leq \liminf \mathbb{E}Z_t^{T_n} = 1,$$

so

$$\mathbb{E}Z_t \leq 1 \quad \forall t. \quad (*)$$

This will be used as an upper bound in an application of the dominated convergence theorem. By application of Itô's formula to

$$f(x) = \frac{x}{1 + \epsilon x},$$

we obtain that

$$f(Z_t) = f(Z_0) + \int_0^t f'(Z_s) dZ_s + \frac{1}{2} \int_0^t f''(Z_s) d\langle Z_s, Z_s \rangle,$$

Hence

$$\frac{Z_t}{1 + \epsilon Z_t} = \frac{1}{1 + \epsilon} - \int_0^t \frac{Z_s h^T(s, X_s)}{(1 + \epsilon Z_s)^2} dW_s - \int_0^t \frac{\epsilon Z_s^2 |h(s, X_s)|^2}{(1 + \epsilon Z_s)^3} ds. \quad (**)$$

Consider the term

$$\int_0^t \frac{Z_s h^T(s, X_s)}{(1 + \epsilon Z_s)^2} dW_s,$$

clearly this a local martingale, since it is a stochastic integral. The next step in the proof is explained in detail as it is one which occurs frequently. We wish to show that the above stochastic integral is in fact a genuine martingale. From the earlier theory (for  $L^2$  integrable martingales) it suffices to show that integrand is in  $L^2(W)$ . We therefore compute

$$\left\| \frac{Z_s h^T(s, X_s)}{(1 + \epsilon Z_s)^2} \right\|_W = \mathbb{E} \left[ \int_0^t \left\| \frac{Z_s h^T(s, X_s)}{(1 + \epsilon Z_s)^2} \right\|^2 ds \right],$$

with a view to showing that it is finite, whence the integrand must be in  $L^2(W)$  and hence the integral is a genuine martingale. We note that since  $Z_s \geq 0$ , that

$$\frac{1}{(1 + \epsilon Z_s)} \leq 1, \text{ and } \frac{Z_s^2}{(1 + \epsilon Z_s)^2} \leq \frac{1}{\epsilon^2},$$

so,

$$\begin{aligned} \left\| \frac{Z_s h(s, X_s)}{(1 + \epsilon Z_s)^2} \right\|^2 &= \frac{Z_s^2}{(1 + \epsilon Z_s)^4} \|h(s, X_s)\|^2 \\ &= \left[ \frac{Z_s^2}{(1 + \epsilon Z_s)^2} \right] \times \left[ \frac{1}{(1 + \epsilon Z_s)^2} \right] \times \|h(s, X_s)\|^2 \\ &\leq \frac{1}{\epsilon^2} \|h(s, X_s)\|^2. \end{aligned}$$

Using this inequality we obtain

$$\mathbb{E} \left[ \int_0^t \left\| \frac{Z_s h^T(s, X_s)}{(1 + \epsilon Z_s)^2} \right\|^2 ds \right] \leq \frac{1}{\epsilon^2} \mathbb{E} \left[ \int_0^T \|h(s, X_s)\|^2 ds \right],$$

however as a consequence of our linear growth condition on  $h$ , the last expectation is finite. Therefore

$$\int_0^t \frac{Z_s h^T(s, X_s)}{(1 + \epsilon Z_s)^2} dW_s$$

is a genuine martingale, and hence taking the expectation of (\*\*) we obtain

$$\mathbb{E} \left[ \frac{Z_t}{1 + \epsilon Z_t} \right] = \frac{1}{1 + \epsilon} - \mathbb{E} \left[ \int_0^t \frac{\epsilon Z_s^2 |h(s, X_s)|^2}{(1 + \epsilon Z_s)^3} ds \right].$$

Consider the integrand on the right hand side; clearly it tends to zero as  $\epsilon \rightarrow 0$ . But also

$$\begin{aligned} \frac{\epsilon Z_s^2 |h(s, X_s)|^2}{(1 + \epsilon Z_s)^3} &= \frac{\epsilon Z_s}{(1 + \epsilon Z_s)^3} \cdot Z_s \cdot |h(s, X_s)|^2 \\ &\leq \frac{1 + \epsilon Z_s}{(1 + \epsilon Z_s)^3} \cdot Z_s \cdot |h(s, X_s)|^2 \\ &\leq k Z_s (1 + \|X_s\|^2), \end{aligned}$$

where we have used the fact that  $|h(s, X_s)|^2 \leq k(1 + \|X_s\|^2)$ . So from the fact that  $\mathbb{E}Z_s \leq 1$  and using the next lemma we shall see that  $\mathbb{E}(Z_s \|X_s\|^2) \leq C$ . Hence we conclude that  $kZ_s(1 + \|X_s\|^2)$  is an integrable dominating function for the integrand on interest. Hence by the Dominated Convergence Theorem, as  $\epsilon \rightarrow 0$ ,

$$\mathbb{E} \left[ \int_0^t \frac{\epsilon Z_s^2 |h(s, X_s)|^2}{(1 + \epsilon Z_s)^3} ds \right] \rightarrow 0.$$

In addition, since  $\mathbb{E}(Z_s) \leq 1$ , we have

$$\mathbb{E} \left( \frac{Z_s}{1 + \epsilon Z_s} \right) \rightarrow \mathbb{E}(Z_s), \text{ as } \epsilon \rightarrow 0.$$

Hence we conclude that

$$\mathbb{E}(Z_t) = 1 \quad \forall t,$$

and so  $Z_t$  is a genuine martingale. □

**Lemma (Gronwall's Inequality).**

Suppose that  $g(t)$  is a continuous function which satisfies

$$0 \leq g(t) \leq \alpha(t) + \beta \int_0^t g(s) ds; \quad 0 \leq t \leq T,$$

in which  $\beta \geq 0$  and  $\alpha : [0, T] \rightarrow \mathbb{R}$  is integrable. Then

$$g(t) \leq \alpha(t) + \beta \int_0^t \alpha(s) e^{\beta(t-s)} ds; \quad 0 \leq t \leq T.$$

**Lemma 16.2.**

$$\mathbb{E} \left[ \int_0^t Z_s \|X_s\|^2 ds \right] < C(T) \quad \forall t \in [0, T].$$

*Proof*

To establish this result Itô's formula is used to derive two important results

$$\begin{aligned} d(\|X_t\|^2) &= 2X_t^T (f dt + \sigma dW_s) + \text{tr}(\sigma^T \sigma) dt \\ d(Z_t \|X_t\|^2) &= -Z_t \|X_t\|^2 h^T dV_s + Z_t (2X_t^T (f dt + \sigma dW_s) + \text{tr}(\sigma \sigma^T)) dt \end{aligned}$$

Putting these into the following larger expression yields

$$\begin{aligned} d \left( \frac{Z_t \|X_t\|^2}{1 + \epsilon Z_t \|X_t\|^2} \right) &= \frac{1}{(1 + \epsilon Z_t \|X_t\|^2)^2} d(Z_t \|X_t\|^2) + \\ &\quad \frac{\epsilon}{(1 + \epsilon Z_t \|X_t\|^2)^3} d\langle Z_t \|X_t\|^2, Z_t \|X_t\|^2 \rangle. \end{aligned}$$

Inserting the appropriate expressions for  $d(Z_t \|X_t\|^2)$  yields

$$\begin{aligned} d \left( \frac{Z_t \|X_t\|^2}{1 + \epsilon Z_t \|X_t\|^2} \right) &= \frac{1}{(1 + \epsilon Z_t \|X_t\|^2)^2} [-Z_t \|X_t\|^2 h^T dW_t + Z_t 2X_t^T \sigma dV_s] \\ &\quad + \left[ \frac{Z_t (2X_t^T f + \text{tr}(\sigma \sigma^T))}{(1 + \epsilon Z_t \|X_t\|^2)^2} - \frac{\epsilon (Z_t^2 \|X_t\|^4 h^T h + 4Z_t^2 2X_t^T \sigma \sigma^T X_t)}{(1 + \epsilon Z_t \|X_t\|^2)^3} \right] dt. \end{aligned}$$

After integrating this expression from 0 to  $t$ , the terms which are stochastic integrals are clearly local martingales. In fact they can be shown to be genuine martingales. For example consider the term

$$\int_0^t \frac{Z_s 2X_s^T \sigma}{(1 + \epsilon Z_s \|X_s\|^2)^2} dV_s,$$

we much therefore check that

$$\int_0^t \left[ \frac{Z_s 2X_s^T \sigma}{(1 + \epsilon Z_s \|X_s\|^2)^2} \right]^2 ds = 4 \int_0^t \frac{Z_s^2 X_s^T \sigma \sigma^T X_s}{(1 + \epsilon Z_s \|X_s\|^2)^4} ds < \infty.$$

Note that the term  $X_t^T \sigma \sigma^T X_t$  is a sum over terms of the form

$$|X_t^i \sigma_{ij}(t, X_t) \sigma_{kj}(t, X_t) X_t^k| \leq \|X_t\|^2 \|\sigma\|^2,$$

of which there are  $d^3$  terms (in  $\mathbb{R}^d$ ). But by the linear increase condition on  $\sigma$ , we have for some constant  $\kappa$

$$\|\sigma\|^2 \leq \kappa(1 + \|X\|^2),$$

and hence

$$|X_t^i \sigma_{ij}(t, X_t) \sigma_{kj}(t, X_t) X_t^k| \leq \kappa \|X_t\|^2 (1 + \|X_t\|^2),$$

so the integral may be bounded by

$$\begin{aligned} \int_0^t \frac{Z_s^2 X_s^T \sigma \sigma^T X_s}{(1 + \epsilon Z_s \|X_s\|^2)^4} ds &\leq \kappa d^3 \int_0^t \frac{Z_s^2 \|X_s\|^2 (1 + \|X_s\|^2)}{(1 + \epsilon Z_s \|X_s\|^2)^4} ds \\ &= \kappa d^3 \left[ \int_0^t \frac{Z_s^2 \|X_s\|^2}{(1 + \epsilon Z_s \|X_s\|^2)^4} ds + \int_0^t \frac{Z_s^2 \|X_s\|^4}{(1 + \epsilon Z_s \|X_s\|^2)^4} ds \right] \end{aligned}$$

Considering each term separately,

$$\begin{aligned} \int_0^t \frac{Z_s^2 \|X_s\|^2}{(1 + \epsilon Z_s \|X_s\|^2)^4} ds &\leq \int_0^t Z_s \times \frac{Z_s \|X_s\|^2}{(1 + \epsilon Z_s \|X_s\|^2)} \times \frac{1}{(1 + \epsilon Z_s \|X_s\|^2)^3} ds \\ &\leq \int_0^t \frac{Z_s}{\epsilon} ds \leq \frac{1}{\epsilon} \int_0^t Z_s, \end{aligned}$$

the last integral is bounded because  $\mathbb{E}(Z_s) \leq 1$ . Similarly for the second term,

$$\int_0^t \frac{Z_s^2 \|X_s\|^4}{(1 + \epsilon Z_s \|X_s\|^2)^4} ds \leq \int_0^t \frac{Z_s^2 \|X_s\|^4}{(1 + \epsilon Z_s \|X_s\|^2)^2} \times \frac{1}{(1 + \epsilon Z_s \|X_s\|^2)^2} ds \leq \frac{1}{\epsilon^2} t < \infty.$$

A similar argument holds for the other stochastic integral, so they they must both be genuine martingales, and hence if we take the expectations of the integrals they are zero. Hence integrating the whole expression from 0 to  $t$ , the taking expectation and finally differentiation with respect to  $t$  yields

$$\begin{aligned} \frac{d}{dt} \mathbb{E} \left( \frac{Z_t \|X_t\|^2}{1 + \epsilon Z_t \|X_t\|^2} \right) &\leq \mathbb{E} \left( \frac{Z_t (2X_t^T f + \text{tr}(\sigma \sigma^T))}{(1 + \epsilon Z_t \|X_t\|^2)^2} \right) \\ &\leq k \left( 1 + \mathbb{E} \left[ \frac{Z_t \|X_t\|^2}{1 + \epsilon Z_t \|X_t\|^2} \right] \right). \end{aligned}$$

Applying Gronwall's inequality gives,

$$\mathbb{E} \left( \frac{Z_t \|X_t\|^2}{1 + \epsilon Z_t \|X_t\|^2} \right) \leq C(T); \quad \forall t \in [0, T].$$

Applying Fatou's lemma the desired result is obtained.  $\square$

Given that  $Z_t$  has been shown to be a martingale a new probability measure  $\tilde{\mathbb{P}}$  may be defined by the Radon-Nikodym derivative:

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} \Big|_{\mathcal{F}_T} = Z_T.$$

**Proposition 16.3.**

Under the new measure  $\tilde{\mathbb{P}}$ ,  $Y$  is a Brownian motion, independent of  $X$ .

*Proof*

From Girsanov's theorem, the process

$$Y_t = V_t + \int_0^t h(s, X_s) ds,$$

is a Brownian Motion under the new measure  $\tilde{\mathbb{P}}$ . Now we must check independence of  $Y$  and  $X$ . The law of the pair  $(X, Y)$  on  $[0, T]$  is absolutely continuous with respect to that of the pair  $(X, V)$  on  $[0, T]$ , since the latter is equal to the former plus a drift term.

Their Radon-Nikodym derivative (which exists since they are absolutely continuous with respect to each other) is  $Z_T$ , which means that for  $f$  any bounded measurable function

$$\mathbb{E}_{\tilde{\mathbb{P}}} [f(X, Y) Z_T] = \mathbb{E}_{\mathbb{P}} [f(X, V)].$$

Hence in terms of the new measure

$$\mathbb{E}_{\tilde{\mathbb{P}}} [f(X, Y)] = \mathbb{E}_{\mathbb{P}} [f(X, V)],$$

and hence under  $\tilde{\mathbb{P}}$ , the variables  $X$  and  $Y$  are independent.  $\square$

**Proposition 16.4.**

Let  $U$  be an integrable  $\mathcal{F}_t$  measurable random variable. Then

$$\mathbb{E}_{\tilde{\mathbb{P}}} [U | \mathcal{Y}_t] = \mathbb{E}_{\tilde{\mathbb{P}}} [U | \mathcal{Y}].$$

*Proof*

Define

$$\mathcal{Y}'_t := \sigma(Y_{t+u} - T_t : u \geq 0),$$

then  $\mathcal{Y} = \mathcal{Y}_t \cup \mathcal{Y}'_t$ . Under the measure  $\tilde{\mathbb{P}}$ , the  $\sigma$ -algebra  $\mathcal{Y}'_t$  is independent of  $\mathcal{F}_t$ , since  $Y$  is an  $\mathcal{F}_t$  adapted Brownian Motion. Hence

$$\mathbb{E}_{\tilde{\mathbb{P}}} [U | \mathcal{Y}_t] = \mathbb{E}_{\tilde{\mathbb{P}}} [U | \mathcal{Y}_t \cup \mathcal{Y}'_t] = \mathbb{E}_{\tilde{\mathbb{P}}} [U | \mathcal{Y}].$$

$\square$

**16.5. The Unnormalised Conditional Distribution**

To simplify the notation in what follows define  $\tilde{Z}_t = 1/Z_t$ . Then under the measure  $\tilde{\mathbb{P}}$  we have

$$d\tilde{Z}_t = \tilde{Z}_t h^T(t, X_t) dY_t, \quad (*)$$

so  $\tilde{Z}_t$  is a  $\tilde{\mathbb{P}}$  local martingale. Hence

$$\tilde{Z}_t = \exp \left( \int_0^t h^T(s, X_s) dY_t - \frac{1}{2} \int_0^t |h(s, X_s)|^2 ds \right).$$

Also  $\mathbb{E}_{\tilde{\mathbb{P}}}(\tilde{Z}_t) = \mathbb{E}_{\mathbb{P}}(Z_t \tilde{Z}_t) = 1$ , the constant mean condition implies that  $\tilde{Z}_t$  is a  $\tilde{\mathbb{P}}$  martingale. Also

$$\frac{d\tilde{\mathbb{P}}}{d\mathbb{P}} \Big|_{\mathcal{F}_t} = \tilde{Z}_t, \text{ for all } t \geq 0.$$

Now **define**, for every bounded measurable function  $\phi$ , the *unnormalised conditional distribution on  $X$*  via

$$\begin{aligned} \rho_t(\phi) &:= \mathbb{E}_{\tilde{\mathbb{P}}} \left[ \phi(X_t) \tilde{Z}_t | \mathcal{Y}_t \right] \\ &= \mathbb{E}_{\tilde{\mathbb{P}}} \left[ \phi(X_t) \tilde{Z}_t | \mathcal{Y} \right], \end{aligned}$$

using the result of proposition 16.4. □

**Proposition (Kallianpur-Striebel) 16.5.**

For every bounded measurable function  $\phi$ , we have

$$\pi_t(\phi) := \mathbb{E}_{\mathbb{P}} [\phi(X_t) | \mathcal{Y}_t] = \frac{\rho_t(\phi)}{\rho_t(1)}, \text{ } \mathbb{P} \text{ and } \tilde{\mathbb{P}} \text{ a.s..}$$

*Proof*

Let  $b$  be a bounded  $\mathcal{Y}_t$  measurable function. From the definition that

$$\pi_t(\phi) := \mathbb{E}_{\mathbb{P}} [\phi(X_t) | \mathcal{Y}_t],$$

we deduce from the definition of conditional expectation that

$$\mathbb{E}_{\mathbb{P}} (\pi_t(\phi) b) = \mathbb{E}_{\mathbb{P}} (\phi(X_t) b),$$

hence

$$\mathbb{E}_{\tilde{\mathbb{P}}} (\tilde{Z}_t \pi_t(\phi) b) = \mathbb{E}_{\tilde{\mathbb{P}}} (\tilde{Z}_t \phi(X_t) b).$$

Whence

$$\mathbb{E}_{\tilde{\mathbb{P}}} (\pi_t(\phi) \mathbb{E}_{\tilde{\mathbb{P}}} [\tilde{Z}_t | \mathcal{Y}_t] b) = \mathbb{E}_{\tilde{\mathbb{P}}} (\mathbb{E}_{\tilde{\mathbb{P}}} [\phi(X_t) \tilde{Z}_t | \mathcal{Y}_t] b).$$

□

## 16.6. The Zakai Equation

**Definition 6. Definition**

A set  $S$  is said to be *total* in  $L^1(\Omega, \mathcal{F}_t, \mathbb{P})$ , if for every  $a \in L^1(\Omega, \mathcal{F}_t, \mathbb{P})$

$$\mathbb{E}(a \epsilon_t) = 0 \quad \forall \epsilon \in S \Rightarrow a = 0.$$

**Lemma 16.7.**

Let

$$S_t := \left\{ \epsilon_t = \exp \left( i \int_0^t r_s^T dY_s + \frac{1}{2} \int_0^t r_s^T r_s ds \right) \quad : \quad r_s \in L^\infty([0, t], \mathbb{R}^m) \right\},$$

then the set  $S_t$  is a total set in  $L^1(\Omega, \mathcal{Y}_t, \tilde{\mathbb{P}})$ .

**Remark**

As a result if  $\phi \in L^1(\Omega, \mathcal{Y}_t, \tilde{\mathbb{P}})$  and we wish to compute  $\mathbb{E}(\phi|\mathcal{Y})$ , it suffices to compute  $\mathbb{E}(\phi\epsilon_t)$  for all  $\epsilon_t \in S_t$ .

**Lemma 16.8.**

Let  $\{U_t\}_{t \geq 0}$  be a continuous  $\mathcal{F}_t$  measurable process such that

$$\tilde{\mathbb{E}} \left( \int_0^T U_t^2 dt \right) < \infty, \quad \forall T \geq 0,$$

then for any  $t \geq 0$ , and for  $j = 1, \dots, m$  the following holds

$$\tilde{\mathbb{E}} \left( \int_0^t U_s dY_s^j | \mathcal{Y}_t \right) = \int_0^t \tilde{\mathbb{E}}(U_s | \mathcal{Y}_s) dY_s^j.$$

*Proof*

As  $S_t$  is a total set, it is sufficient to prove that for any  $\epsilon_t \in S_t$  that

$$\tilde{\mathbb{E}} \left( \epsilon_t \tilde{\mathbb{E}} \left( \int_0^t U_s dY_s^j | Y_t \right) \right) = \tilde{\mathbb{E}} \left( \epsilon_t \tilde{\mathbb{E}}(U_s | Y_s) dY_s^j \right).$$

To this end note that for every  $\epsilon_t \in S_t$  the following holds

$$\epsilon_t = 1 + \int_0^t i\epsilon_s r_s^T dY_s,$$

and hence

$$\begin{aligned} \tilde{\mathbb{E}} \left( \epsilon_t \tilde{\mathbb{E}} \left( \int_0^t U_s dY_s^j | Y_t \right) \right) &= \tilde{\mathbb{E}} \left( \epsilon_t \int_0^t U_s dY_s^j \right) \\ &= \tilde{\mathbb{E}} \left( \left( 1 + \int_0^t i\epsilon_s r_s^T dY_s \right) \left( \int_0^t U_s dY_s^j \right) \right) \\ &= \tilde{\mathbb{E}} \left( \int_0^t U_s dY_s^j \right) + \tilde{\mathbb{E}} \left( \int_0^t i\epsilon_s r_s^j U_s ds \right). \end{aligned}$$

Here the last term is computed by recalling the definition of the covariation process, in conjunction with the Kunita Watanabe identity. The first term is a martingale (from the boundedness condition on  $U_s$ ) and hence its expectation vanishes. Thus

$$\begin{aligned} \tilde{\mathbb{E}} \left( \epsilon_t \tilde{\mathbb{E}} \left( \int_0^t U_s dY_s^j | Y_t \right) \right) &= \tilde{\mathbb{E}} \left( \tilde{\mathbb{E}} \left( \int_0^t i\epsilon_s r_s^j U_s ds | \mathcal{Y}_t \right) \right) \\ &= \tilde{\mathbb{E}} \left( \int_0^t i\epsilon_s r_s^j \tilde{\mathbb{E}}(U_s | \mathcal{Y}_s) ds \right) \\ &= \tilde{\mathbb{E}} \left( \epsilon_t \int_0^t \tilde{\mathbb{E}}(U_s | \mathcal{Y}_s) dY_s^j \right). \end{aligned}$$

As this holds for all  $\epsilon_t$  in  $S_t$ , and the latter is a total set, by the earlier remarks, this establishes the result.  $\square$

**Lemma 16.9.**

Let  $\{U_t\}_{t \geq 0}$  be an  $\mathcal{F}_t$  adapted continuous process such that

$$\tilde{\mathbb{E}} \left( \int_0^T U_s^2 ds \right) < \infty, \quad \forall T \geq 0, \quad (*)$$

and let  $R_t$  be another continuous  $\mathcal{F}_t$  adapted process such that  $\langle R, Y \rangle_t = 0$  for all  $t$ . Then

$$\tilde{\mathbb{E}} \left[ \int_0^t U_s dR_s | \mathcal{Y}_t \right] = 0.$$

*Proof*

As before use the fact that  $S_t$  is a total set, so it suffices to show that

$$\tilde{\mathbb{E}} \left( \epsilon_t \tilde{\mathbb{E}} \left( \int_0^t U_s dR_s | \mathcal{Y}_t \right) \right) = 0.$$

Now as  $\epsilon_t$  is  $\mathcal{Y}_t$  measurable we see that

$$\begin{aligned} \tilde{\mathbb{E}} \left( \epsilon_t \tilde{\mathbb{E}} \left( \int_0^t U_s dR_s | \mathcal{Y}_t \right) \right) &= \tilde{\mathbb{E}} \left( \tilde{\mathbb{E}} \left( \epsilon_t \int_0^t U_s dR_s | \mathcal{Y}_t \right) \right) \\ &= \tilde{\mathbb{E}} \left( \epsilon_t \int_0^t U_s dR_s \right) \end{aligned}$$

However as before each  $\epsilon_t$  in  $S_t$  satisfied

$$\epsilon_t = 1 + \int_0^t i \epsilon_s r_s^T dY_s$$

Hence substituting this into the above expression yields

$$\begin{aligned} \tilde{\mathbb{E}} \left( \epsilon_t \tilde{\mathbb{E}} \left( \int_0^t U_s dR_s | \mathcal{Y}_t \right) \right) &= \tilde{\mathbb{E}} \left( \int_0^t U_s dR_s \right) + \tilde{\mathbb{E}} \left[ \left( \int_0^t i \epsilon_s r_s^T dY_s \right) \left( \int_0^t U_s dR_s \right) \right] \\ &= \tilde{\mathbb{E}} \left( \int_0^t U_s dR_s \right) + \tilde{\mathbb{E}} \left( \int_0^t i \epsilon_s U_s r_s^T d\langle Y, R \rangle \right) \\ &= \tilde{\mathbb{E}} \left( \int_0^t U_s dR_s \right) \\ &= 0. \end{aligned}$$

The last term vanishes, because by the condition (\*), the stochastic integral is a genuine martingale. The other term vanishes because from the hypotheses  $\langle Y, R \rangle = 0$ .  $\square$

**Remark**

In the context of stochastic filtering a natural application of this lemma will be made by setting  $R_t = V_t$ , the process driving the signal process.

We are now in a position to state and prove the Zakai equation. The Zakai equation is important because it is a parabolic stochastic partial differential equation which is satisfied by  $\rho_t$ , and indeed its solution provides a practical means of computing  $\rho_t(\phi) = \tilde{\mathbb{E}}(\phi(X_t)\tilde{Z}_t|\mathcal{Y}_t)$ . Indeed the Zakai equation provides a method to approach numerical solution of the non-linear filtering problem by using recursive algorithms to solve the stochastic PDE.

**Theorem 16.10.**

Let  $A$  be the infinitesimal generator of the signal process  $X_t$ , let the domain of this infinitesimal generator be denoted  $\mathcal{D}(A)$ . Then subject to the usual conditions on  $f$ ,  $\sigma$  and  $h$ , the un-normalised conditional distribution on  $X$  satisfies the Zakai equation which is

$$\rho_t(\phi) = \pi_0(\phi) + \int_0^t \rho_s(A_s\phi)ds + \int_0^t \rho_s(h^T\phi)dY_s, \quad \forall t \geq 0, \quad \forall \phi \in \mathcal{D}(A).$$

*Proof*

We approximate  $\tilde{Z}_t$  by

$$\tilde{Z}_t^\epsilon = \frac{\tilde{Z}_t}{1 + \epsilon\tilde{Z}_t},$$

from the integration by parts formula,

$$d(\tilde{Z}_t^\epsilon\phi(X_t)) = \phi(X_t)d\tilde{Z}_t^\epsilon + \tilde{Z}_t^\epsilon d(\phi(X_t)) + \langle \tilde{Z}_t^\epsilon, \phi(X_t) \rangle.$$

However from the definition of an infinitesimal generator, we know that

$$\phi(X_t) - \phi(X_0) = \int_0^t A_s\phi(x)ds + M_t^\phi,$$

where  $M_t^\phi$  is a martingale. Also application of Itô's formula to  $f(x) = x/(1 + \epsilon x)$ , yields

$$d\tilde{Z}_t^\epsilon = df(\tilde{Z}_t) = (1 + \epsilon\tilde{Z}_t)^{-2}\tilde{Z}_t h^T dY_t - \epsilon(1 + \epsilon\tilde{Z}_t)^{-3}\tilde{Z}_t^2 \|h(t, X_t)\|^2.$$

Putting these two results together yields

$$\begin{aligned} d(\tilde{Z}_t^\epsilon\phi(X_t)) &= \left[ \tilde{Z}_t^\epsilon A\phi(X_t) - \epsilon(1 + \epsilon\tilde{Z}_t)^{-3}\|h(t, X_t)\|^2 \right] dt \\ &\quad + \tilde{Z}_t^\epsilon dM_t^\phi + \phi(X_t)(1 + \epsilon\tilde{Z}_t)^{-2}\tilde{Z}_t h^T(t, X_t)dY_t \end{aligned}$$

In terms of the functions already defined we can write the infinitesimal generator for  $X_t$  as

$$A\phi(x) = \sum_{i=1}^d f_i(s, x) \frac{\partial \phi}{\partial x^i}(x) + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d \left( \sum_{k=1}^d \sigma_{ik}(s, x) \sigma_{kj}(s, x) \right) \frac{\partial^2 \phi}{\partial x^i \partial x^j}(x),$$

and from the earlier discussion of the martingale problem for  $A$ , we can also write

$$M_t^\phi = \int_0^t \sum_{i=1}^d \sum_{j=1}^d \frac{\partial \phi}{\partial x^i}(X_t) \sigma_{ij}(t, X_t) dW_t^j.$$

Now we wish to compute

$$\tilde{\mathbb{E}} \left( \tilde{Z}_t^\epsilon \phi(X_t) | \mathcal{Y}_t \right)$$

we shall do this using the previous expression which on integration gives us

$$\begin{aligned} \tilde{Z}_t^\epsilon \phi(X_t) &= \tilde{Z}_0^\epsilon \phi(X_0) + \int_0^t \left[ \tilde{Z}_s^\epsilon A \phi(X_s) - \epsilon (1 + \epsilon \tilde{Z}_s) \|\tilde{h}(s, X_s)\|^2 \right] ds \\ &\quad + \int_0^t \tilde{Z}_s^\epsilon dM_s^\phi + \int_0^t \phi(X_s) (1 + \epsilon \tilde{Z}_s)^{-2} \tilde{Z}_s h^T(s, X_s) dY_s \end{aligned}$$

Note that

$$\tilde{\mathbb{E}} \left( \tilde{Z}_0^\epsilon \phi(X_0) | \mathcal{Y}_t \right) = \rho_0(\phi) = \pi_0(\phi) \rho_0(1) = \pi_0(\phi) \frac{1}{1 + \epsilon}.$$

Hence taking conditional expectations on the whole expression we obtain

$$\begin{aligned} \tilde{\mathbb{E}} \left( \tilde{Z}_t^\epsilon \phi(X_t) | \mathcal{Y}_t \right) &= \frac{\pi_0(\phi)}{1 + \epsilon} + \int_0^t \tilde{\mathbb{E}} \left( \phi(X_s) (1 + \epsilon \tilde{Z}_s)^{-2} \tilde{Z}_s h^T(s, X_s) | \mathcal{Y}_t \right) dY_s \\ &\quad + \dots \end{aligned}$$

Now we take the  $\epsilon \downarrow 0$  limit. Clearly  $\tilde{Z}_t^\epsilon \rightarrow \tilde{Z}_t$ , and also

$$\tilde{\mathbb{E}} \left( \tilde{Z}_t^\epsilon \phi(X_t) | \mathcal{Y}_t \right) \rightarrow \tilde{\mathbb{E}} \left( \tilde{Z}_t \phi(X_t) | \mathcal{Y}_t \right) = \rho_t(\phi),$$

and also

$$\tilde{\mathbb{E}} \left( \tilde{Z}_s^\epsilon A_s \phi(X_s) | \mathcal{Y}_s \right) \rightarrow \rho_s(A_s \phi).$$

We can bound this by

$$\tilde{\mathbb{E}} \left( \tilde{Z}_s^\epsilon A_s \phi(X_s) | \mathcal{Y}_s \right) \leq \|A_s \phi\| \tilde{\mathbb{E}} \left( \tilde{Z}_s | \mathcal{Y}_s \right),$$

and the right hand side is an  $L^1$  bounded quantity. Hence by the dominated convergence theorem we have

$$\int_0^t \tilde{\mathbb{E}} \left( \tilde{Z}_s^\epsilon A_s \phi(X_s) | \mathcal{Y}_s \right) ds \rightarrow \int_0^t \rho_s(A_s \phi) ds \text{ a.s.}$$

Also note that

$$\lim_{\epsilon \rightarrow 0} \epsilon \phi(X_s) \left( \tilde{Z}_s^\epsilon \right)^2 \left( 1 + \epsilon \tilde{Z}_s \right)^{-1} \|\tilde{h}(s, X_s)\|^2 = 0.$$

and since the function may be dominated by

$$\left| \epsilon \phi(X_s) (Z_s^\epsilon)^2 (1 + \tilde{Z}_s)^{-1} \|h(s, X_s)\|^2 \right| \leq \|\phi\| \tilde{Z}_s \|h(s, X_s)\|^2,$$

as a consequence of these two results, we see via the Dominated Convergence theorem, as the right hand side of the above inequality is integrable, that

$$\int_0^t \epsilon \tilde{\mathbb{E}} \left[ \phi(X_s) (\tilde{Z}_s^\epsilon)^2 (1 + \epsilon \tilde{Z}_s)^{-1} \|h(s, X_s)\|^2 | \mathcal{Y}_s \right] ds \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0.$$

It now remains to check that the stochastic integral

$$\int_0^t \tilde{\mathbb{E}} \left[ \phi(X_s) \tilde{Z}_s^\epsilon (1 + \tilde{Z}_s)^{-1} h^T(s, X_s) | \mathcal{Y}_s \right] dY_s \rightarrow \int_0^t \rho_s(h^T \phi(X_s)) dY_s.$$

To this end define

$$I_\epsilon(t) := \int_0^t \tilde{\mathbb{E}} \left[ \phi(X_s) \tilde{Z}_s^\epsilon (1 + \tilde{Z}_s)^{-1} h^T(s, X_s) | \mathcal{Y}_s \right] dY_s,$$

and the desired limit

$$I(t) := \int_0^t \rho_s(h^T \phi(X_s)) dY_s.$$

Note that both of these can be shown to be continuous, square bounded integrable martingales. Now note that the difference

$$I_\epsilon(t) - I(t) = - \int_0^t \tilde{\mathbb{E}} \left[ \phi(X_s) \frac{\epsilon \tilde{Z}_s^2 (2 + \epsilon \tilde{Z}_s)}{(1 + \epsilon \tilde{Z}_s)^2} h^T(s, X_s) | \mathcal{Y}_s \right] dY_s,$$

Now we use the dominated convergence theorem to show that for a suitable subsequence that since  $\|h(s, X_s)\|^2 \leq K(1 + \|X_s\|^2)$  we can write

$$\tilde{\mathbb{E}} \left[ \int_0^t \left( \tilde{\mathbb{E}} \left[ \phi(X_s) \frac{\epsilon \tilde{Z}_s^2 (2 + \epsilon \tilde{Z}_s)}{(1 + \epsilon \tilde{Z}_s)^2} h^T(s, X_s) | \mathcal{Y}_s \right] \right)^2 ds \right] \rightarrow 0. \quad a.s.$$

which is the same as

$$\tilde{\mathbb{E}} \left[ (I_\epsilon(t) - I(t))^2 \right] \rightarrow 0.$$

By a standard theorem of convergence, this means that there is a suitable subsequence  $\epsilon_n$  such that

$$I_{\epsilon_n}(t) \rightarrow I(t) \quad a.s.$$

which is sufficient to establish the claim.  $\square$

### 16.7. Kushner-Stratonovich Equation

The Zakai equation as described above provides an SDE satisfied by the unnormalised conditional distribution of  $X$ , i.e.  $\rho_t(\phi)$ . It seems natural therefore to derive a similar equation satisfied by the normalised conditional distribution,  $\pi_t(\phi)$ . Recall that the unnormalised conditional distribution is given by

$$\rho_t(\phi) = \tilde{\mathbb{E}} \left[ \phi(X_t) \tilde{Z}_t | \mathcal{Y}_t \right],$$

and the normalised conditional law is given by

$$\pi_t(\phi) := \mathbb{E} [\phi(X_t) | \mathcal{Y}_t].$$

As a consequence of the Kallianpur-Striebel formula

$$\pi_t(\phi) = \frac{\rho_t(\phi)}{\rho_t(1)},$$

Now using this result together with the Zakai equation, we can derive the Kushner-Stratonovich equation.

**Theorem (Kushner-Stratonovich) 16.11.**

*The normalised conditional law of the process  $X_t$  subject to the usual conditions on the processes  $X$  and  $Y$  satisfies the Kushner-Stratonovich equation*

$$\pi_t(\phi) = \pi_0(\phi) + \int_0^t \pi_s(A_s \phi) ds + \int_0^t (\pi_s(h^T \phi) - \pi_s(h^T) \pi_s(\phi)) (dY_s - \pi_s(h) ds).$$